

Molecular Neurobiology for Practicing Psychiatrists, Part 5: How a Leucine Zipper Can Turn On Genes: Immediate-Early Genes Activate Late-Genes Expression in the Brain

Stephen M. Stahl, M.D., Ph.D.

Issue: *Neurotransmitters activate genes that activate other genes; that is, neurotransmission can rapidly deploy a strike force of immediate-early genes, like the first troops sent into combat. These early genes then combine to draft an army of genes to mobilize numerous important gene products that alter the function of a neuron targeted by a neurotransmitter.*

T

his is the fifth and final lesson in our series explaining molecular neurobiology for psychiatrists (see references 1–5).

Previous lessons in this series have shown how neurotransmitters can quickly activate genes like *c-fos* and *c-jun* (Figure 1).^{3,4} These genes function as rapid responders to the neurotransmitter's input, like the first troops sent into combat once war has been declared. This rapid deployment force of immediate-early genes (IEGs) reacts within 15 minutes of receiving a neurotransmission (Figure 2) by being the first to encode their respective proteins, Fos and Jun. This encoding lasts for only a half hour to an hour, but it is enough time for Fos and Jun, nuclear proteins that live and work in the neuron's cell nucleus, to team up and form a leucine zipper-type of transcription factor. The zipper, in turn, activates many kinds of late-onset genes (Figures 1 and 2). Thus, Fos and Jun serve to wake up the much larger army of inactive genes. Which individual soldier genes are so drafted to active gene duty depends on a number of factors: which neurotransmitter is sending the message,

how frequently it is sending the message, and whether it is working in concert with or opposition to other neurotransmitters talking to other parts of the same neuron at the same time.

REFERENCES

1. Stahl SM. Molecular neurobiology for practicing psychiatrists, part 1: overview of gene activation by neurotransmitters [BRAINSTORMS]. *J Clin Psychiatry* 1999;60:572–573
2. Stahl SM. Molecular neurobiology for practicing psychiatrists, part 2: how neurotransmitters activate second messenger systems [BRAINSTORMS]. *J Clin Psychiatry* 1999;60:647–648
3. Stahl SM. Molecular neurobiology for practicing psychiatrists, part 3: How second messengers "turn on" genes by activating protein kinases and transcription factors. [BRAINSTORMS]. *J Clin Psychiatry* 1999;60:731–732
4. Stahl SM. Molecular neurobiology for practicing psychiatrists, part 4 [BRAINSTORMS]. *J Clin Psychiatry* 1999;60:813–814
5. Stahl SM. *Essential Psychopharmacology*. 2nd ed. New York, NY: Cambridge University Press. In press

Take-Home Points

- ◆ Practicing psychiatrists should become conversant with the vocabulary of molecular neurobiology, because this is now the language describing the biological basis and genetics of mental illnesses.
- ◆ The molecular neurobiology of how neuronal genes are activated can also explain the ultimate mechanism of action of numerous psychotropic drugs used in daily practice by psychiatrists.

BRAINSTORMS is a monthly section of *The Journal of Clinical Psychiatry* aimed at providing updates of novel concepts emerging from the neurosciences that have relevance to the practicing psychiatrist.

From the Clinical Neuroscience Research Center in San Diego and the Department of Psychiatry at the University of California San Diego.

Reprint requests to: Stephen M. Stahl, M.D., Ph.D., Editor, *BRAINSTORMS*, 8899 University Center Lane, Suite 130, San Diego, CA 92122.

Figure 1. How Early Genes Activate Late Genes




Figure 2. Time Course of Neurotransmitter-Induced Activation of Genes

