

Independent Actions on Fear Circuits May Lead to Therapeutic Synergy for Anxiety When Combining Serotonergic and GABAergic Agents

Stephen M. Stahl, M.D., Ph.D.

Issue: *Both serotonin and GABA powerfully regulate the neuroanatomical circuit that mediates fear in anxiety disorders. This suggests that use of pharmacotherapies acting on both systems may provide synergistic therapeutic actions.*

In last month's BRAINSTORMS,¹ we discussed the fact that benzodiazepines are still the leading treatments for anxiety disorders, despite their limitations and being relegated to second-line use in modern treatment guidelines. Benzodiazepines act upon GABAergic neurotransmission to enhance neuronal inhibition² and are frequently used along with antidepressants that enhance serotonergic neurotransmission for the treatment of anxiety disorders today.¹ Here we illustrate the neuroanatomical substrate of fear, showing that both GABA and serotonin have inhibitory actions on the output of the fear response from the amygdala.³⁻⁵

Figure 1.
The Amygdala Is the Brain's Panic Button^a






Figure 2. GABA and Serotonin, and Presumably Successful Cognitive-Behavioral Therapy, From Brain Sources Outside the Amygdala Inhibit Amygdala Alarms^a



^aEmotional inputs to the amygdala frequently use the excitatory neurotransmitter glutamate to ring the alarm. However, triggering of the alarm by glutamate can be tempered by both GABA and serotonin.⁶ GABA interneurons in the cortex and hippocampus inhibit emotional input to the amygdala, as do serotonergic nerve terminals from the raphe. Theoretically, cognitive-behavioral therapy enhances inhibitory tone in the cortex by reprogramming the neurons there as they become desensitized and deconditioned to anxiety-provoking triggers.³

Figure 3. GABA and Serotonin From Inside the Amygdala Inhibit Amygdala Alarms^a



^aGABA interneurons and serotonergic nerve terminals are also present directly in the amygdala itself, acting as potential brakes on amygdala output to the fear response. Thus, agents that boost output from either GABA or serotonin neurons each have at least 2 chances to diminish the precipitation of anxiety and fear, as also suggested by recent clinical studies that combine agents from both classes.⁶

REFERENCES

1. Stahl SM. Don't ask, don't tell, but benzodiazepines are still the leading treatments for anxiety disorders. *J Clin Psychiatry* 2002;63:656-657
2. Stahl SM. *Essential Psychopharmacology*. 2nd ed. New York, NY: Cambridge University Press; 2000
3. Gorman JM, Kent JM, Sullivan GM, et al. Neuro-anatomical hypothesis of panic disorder, revised. *Am J Psychiatry* 2000;157:493-505
4. Coplan JD, Lydiard RB. Brain circuits in panic disorder. *Biol Psychiatry* 1998;44:1264-1276
5. Stutzmann GE, LeDoux JE. GABAergic antagonists block the inhibitory effects of serotonin in the lateral amygdala: a mechanism for modulation of sensory inputs to fear conditioning. *J Neurosci* 1999;19:RC8
6. Goddard AW, Brouette T, Almai A, et al. Early coadministration of clonazepam with sertraline for panic disorder. *Arch Gen Psychiatry* 2001;58: 681-686

BRAINSTORMS is a monthly section of The Journal of Clinical Psychiatry aimed at providing updates of novel concepts emerging from the neurosciences that have relevance to the practicing psychiatrist.

From the Neuroscience Education Institute in Carlsbad, Calif., and the Department of Psychiatry at the University of California San Diego.

Reprint requests to: Stephen M. Stahl, M.D., Ph.D., Editor, *BRAINSTORMS*, Neuroscience Education Institute, 5857 Owens Street, Ste. 102, Carlsbad, CA 92009.