

THE PRIMARY CARE COMPANION
TO THE JOURNAL OF CLINICAL PSYCHIATRY
CME ACTIVITY

VOLUME 9 NUMBER 1

CME ARTICLE

**Impact of Bipolar Disorder: Results From a
Screening Study..**



PRETEST AND OBJECTIVES

To receive your credit certificate immediately for free—

Go to
www.PSYCHIATRIST.com
and complete this activity online.

Articles are selected for credit designation on the basis of the CME Institute's assessment of the needs of readers of *The Primary Care Companion*, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. There are no prerequisites for participation in this CME activity.

To obtain credit, please study the designated article and complete the Posttest and Registration Form.

CME Objective

After studying the article by Stang et al., you should be able to:

- Recognize clinical features of bipolar disorder and the impact of the disorder on patients' work and social lives.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc., is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Designation

The CME Institute of Physicians Postgraduate Press, Inc., designates this educational activity for a maximum of 1 *AMA PRA Category 1 Credit(s)*TM. Physicians should only claim credit commensurate with the extent of their participation in the activity.

Date of Original Release/Review

This educational activity is eligible for AMA PRA Category 1 Credit through February 28, 2009. The latest review of this material was January 2007.

This pretest is designed to facilitate your study of the material.

1. **Research has suggested that patients screening positive for bipolar disorder tended to be much less likely to be married and less likely to be employed than those who screened negative, and almost two thirds of those who screened positive had never received a diagnosis of bipolar disorder.**
 - a. True
 - b. False

For Pretest answers and Posttest, see pages 81–82.

Answers to Vol. 6, No. 6 Posttest 2004

1. c 2. b 3. d 4. c 5. a



Impact of Bipolar Disorder: Results From a Screening Study

Paul Stang, Ph.D.; Cathy Frank, M.D.; Marianne Ulcickas Yood, D.Sc., M.P.H.;
Karen Wells, B.S.; and Steven Burch, Ph.D.

Objective: To characterize the extent and impact of bipolar I and II disorders and rapid cycling in a managed care population using both coded diagnostic claims and clinical screening.

Method: The Mood Disorder Questionnaire (MDQ) was used to identify bipolar disorder among patients attending the psychiatry service of a large Midwestern health system. Suicidal ideation screening questions were also asked, along with a brief set of relevant history and medication questions. Patients scoring positive on the MDQ or identified as bipolar-positive according to DSM-IV criteria by the screening clinician were administered the Work and Social Adjustment Scale and an Employment questionnaire. Descriptive statistics were used to summarize results. The study was conducted from July 2004 to November 2004.

Results: Seventy (6.4%) of 1087 patients had bipolar disorder, 59 of whom completed the entire study. For these patients, the mean time with bipolar disorder was 9.3 (SD 10.2) years. The mean length of the current episode was 10.4 (SD 14.4) weeks, with 22% of patients experiencing a mixed episode, 5% manic-predominant, 12% hypomanic-predominant, and 46% a depressive-predominant episode. Twenty-four percent of patients with bipolar disorder were rapid cycling at the time of their visit; for 5 of these patients, rapid cycling was thought to be related to antidepressant use. Sixty-one percent of patients with bipolar disorder were taking an antidepressant; 69.5% were taking a mood stabilizer. Of these patients with bipolar disorder, 19% were evaluated as high suicidality risk, while 47% were considered moderate risk. Bipolar disorder patients reported problems with employment/employability and social adjustment. About one quarter of these patients ranked problems with family and relationships as marked or severe. Fifty percent of these bipolar disorder patients reported missing at least 1 week of work during the past month; 41% reported fearing the loss of their current job due to their emotional state; and 20% reported being fired/laid off during the past 5 years due to their emotional state.

Conclusions: This research documents some of the clinical features and social and labor-force impact of bipolar disorder in a managed care population and adds several dimensions to data published to date. Fully two thirds of our study subjects with bipolar disorder were found to be at substantial risk of suicide, and bipolar disorder patients in this study reported substantial problems with employment/employability and social functioning.

(Prim Care Companion J Clin Psychiatry 2007;9:42-47)

Received Feb. 27, 2006; accepted June 14, 2006. From Galt Associates, Blue Bell, Pa. (Drs. Stang and Yood); College of Health Sciences, West Chester University, West Chester, Pa. (Dr. Stang); the Department of Psychiatry (Dr. Frank) and the Josephine Ford Cancer Center (Dr. Yood and Ms. Wells), Henry Ford Health System, Detroit, Mich.; Epidemiology and Public Health, Yale University School of Medicine, New Haven, Conn. (Dr. Yood); and GlaxoSmithKline, Research Triangle Park, N.C. (Dr. Burch).

This research was funded by GlaxoSmithKline.

In the spirit of full disclosure and in compliance with all ACCME Essential Areas and Policies, the faculty for this CME article were asked to complete a statement regarding all relevant financial relationships between themselves or their spouse/partner and any commercial interest (i.e., any proprietary entity producing health care goods or services consumed by, or used on, patients) occurring within at least 12 months prior to joining this activity. The CME Institute has resolved any conflicts of interest that were identified. The disclosures are as follows: Dr. Stang has been a consultant to and received grant/research support from GlaxoSmithKline and other pharmaceutical companies. Dr. Burch is an employee of GlaxoSmithKline. Drs. Frank and Yood and Ms. Wells have no personal affiliations or financial relationships with any proprietary entity producing health care goods or services consumed by, or used on, patients to disclose relative to the article.

Corresponding author and reprints: Paul Stang, Ph.D., Galt Associates, Inc., 620 Sentry Parkway, Suite 100, Blue Bell, PA 19422 (e-mail: pstang@drugsafety.com).

Bipolar disorder is one of the world's 10 most disabling conditions, and it affects people across nationality, race, and socioeconomic status.¹ While traditionally defined bipolar disorder (bipolar I) has a prevalence of approximately 1% across all populations, the prevalence of bipolar II disorder, involving episodes of less severe hypomania and major depression, along with bipolar spectrum subtypes, brings the estimated prevalence of all bipolar disorders to over 3% of people in the United States.¹

A 2005 primary care practice-based study² reported that the prevalence of positive screening results for lifetime bipolar disorder was 9.8%, and it did not differ significantly by age, sex, or race/ethnicity. Of those who screened positive for bipolar disorder, 72.3% sought professional help for their symptoms, but only 8.4% reported receiving a diagnosis of bipolar disorder. Similarly, among those who screened positive for bipolar disorder, 68.2% had a current major depressive episode or an anxiety or substance use disorder, and only 6.5% reported taking a mood-stabilizing agent in the past month. In that study, primary care physicians recorded evidence of current depression in 49.0% of those screening positive for bipolar disorder, but did not record a bipolar disorder diagnosis in either administrative billing or the medical record of

TAKE-HOME POINTS

- ◆ Increased screening for bipolar disorder with a diagnostic tool (like the Mood Disorder Questionnaire) is an important step in identifying those patients in need of intervention
- ◆ Two thirds of the patients with bipolar disorder were at substantial risk for suicide
- ◆ Among patients with bipolar disorder, 42% reported that their work was at least markedly affected by their symptoms, and social activities were at least markedly affected in 37%

any of these patients. Health-related quality of life was worse among those with bipolar disorder compared with those who screened negative, as was social and family impact.²

A recent clinic-based study³ of patients taking an antidepressant for depression found that 21.3% screened positive for bipolar disorder symptoms on the Mood Disorder Questionnaire (MDQ). Those screening positive for bipolar disorder symptoms were younger, more likely to be white, more likely to be living alone, much less likely to be married, and less likely to be employed than those who screened negative. Almost two thirds of patients screening positive for bipolar disorder had never received that diagnosis. These findings indicate that bipolar disorder is underdiagnosed and undertreated in primary care, and they suggest some of the serious problems and impairments faced by patients with bipolar disorder.³

In addition to the substantial burden of the symptoms of bipolar disorder itself, a wide range of associated medical problems have been reported in the few studies focused on medical illness in this population, the most common being cardiovascular disease, diabetes mellitus, obesity, and thyroid disease.¹ The coexistence of key medical risk factors related to excessive nicotine use, use of alcohol and other drugs, and co-occurring anxiety disorders and eating disorders may lead to the early onset of medical diseases with poor long-term outcomes.¹

Economic costs of bipolar disorder are also high: a 2001 study⁴ estimated the present value of the lifetime cost of persons with onset of bipolar disorder (in 1998) to be 24 billion U.S. dollars. In that study, average cost per case ranged from \$11,720 for those with a single manic episode to \$624,785 for those with nonresponsive/chronic episodes.⁴

The present study focused on systematically screening attendees of a psychiatry clinic for bipolar disorder and documenting the clinical features and impact that bipolar disorder has on their social and labor-force participation.

METHOD

Study Site

We conducted this study in a large, vertically integrated health system serving the primary and specialty health care needs of Midwestern residents. This system is

affiliated with a multispecialty salaried physician group that provides most of the care for health system patients. The health system owns a large, nonprofit, mixed-model health maintenance organization (HMO). In order to optimize the data available for this study, the population was limited to HMO members. The institutional review board reviewed and approved this research. The study was conducted from July 2004 to November 2004.

Assessment Tools

The MDQ⁵ was utilized as an initial screening tool to identify bipolar disorder among each patient attending the health system psychiatry service. Our target was to screen 1000 consecutive clinic attendees. An MDQ screening score of 7 or more is reported to yield good sensitivity (0.73) and very good specificity (0.90) in the diagnosis of bipolar disorder.⁵ The MDQ was immediately scored by a mental health professional, and, regardless of score, patients were immediately evaluated for bipolar disorder using the DSM-IV bipolar disorder criteria. Those meeting the criteria were further evaluated for their rapid-cycling status in addition to any other relevant clinical/mental health disorders. In addition to the MDQ, suicidal ideation screening questions (developed and implemented by the health system with a working group using data culled from key literature sources^{6,7}; please contact author C.F. for further information) were asked by the psychiatric clinician, and a brief set of relevant history and medication questions were completed. Patients scoring positive on the MDQ or found to be bipolar-positive by the clinician were solicited to provide informed consent. For those who consented, the Work and Social Adjustment Scale,⁸ a 9-point Likert scale from 0 (not at all affected) to 8 (very severely affected), and Employment questionnaire (13 questions about employment history, nature of employment, and impact of depressive and/or manic symptoms on employment history; a copy is available on request from author P.S.) were administered.

Data Analysis

Statistical analyses were performed using SAS statistical software (SAS Institute Inc; Cary, N.C.). Descriptive statistics were used to summarize the results of the MDQ screening, the clinician diagnosis, and the findings on the questionnaires.

Table 1. Number of Patients Meeting Bipolar Disorder Criteria According to the Mood Disorder Questionnaire (MDQ) Versus Clinical DSM-IV Assessment^a

		Clinician DSM-IV Appraisal for Bipolar Disorder		Total
		+	-	
MDQ	+	64	57	121
	-	6	960	966
Total		70	1017	1087

^aSensitivity = 0.91 proportion of those with bipolar disorder who have a positive MDQ. Specificity = 0.94 proportion of those without bipolar disorder who have a negative MDQ. Predictive value positive = 0.53 probability that a positive MDQ is truly positive. Predictive value negative = 0.99 probability that a negative MDQ is truly negative.

RESULTS

Screening

There were 1087 MDQs completed by patients attending the psychiatry service. Of these, 121 patients had MDQ-positive results, 64 of whom were found to be bipolar positive by clinician assessment (Table 1). There were 966 patients with MDQ-negative results, of whom 6 were found by the clinician to be bipolar positive. The MDQ did well at identifying those without bipolar disorder (negative predictive value 0.99); however, the MDQ had a relatively high proportion of false-positives, which reduced its positive predictive value to 0.53. Overall, the MDQ identified 91% of those with bipolar disorder (sensitivity) and 94% of those without (specificity).

Therefore, 70 (6.4%) of 1087 patients attending the psychiatry service were found to have bipolar disorder, 59 of whom provided informed consent and completed the entire questionnaire load. The remainder of the results will focus on these 59 patients.

Patient characteristics. Among those patients who were confirmed by the psychiatrist as having bipolar disorder, the mean time with bipolar disorder was 9.3 years (SD 10.2 years). The mean length of their current episode was 10.4 weeks (SD 14.4 weeks), with 22% experiencing a mixed episode, 5% a manic predominant episode, 12% a hypomanic predominant episode, and 46% a depressive predominant episode. Almost 1 in 4 (24%) were thought to be rapid cycling at the time of their visit.

Treatments received. Of the 59 patients with bipolar disorder who completed all of the surveys, 61% were found to be taking an antidepressant; almost half of whom (42%) were taking selective serotonin reuptake inhibitors (SSRIs). Among these 59 patients, 69.5% were taking a mood stabilizer.

The psychiatry clinician felt that in 5 (36%) of the 14 rapid-cycling patients their rapid cycling was related to antidepressant use. Antidepressant monotherapy was more common among non-rapid cyclers (22.7% vs. 7.4%) as was mood stabilizer monotherapy (27.3% vs.

Table 2. Employment Status Among Surveyed Bipolar Disorder Patients

Current Work Situation	N	%
Work at least 40 hours per week for pay	23	38.98
Work 20–39 hours per week for pay	4	6.78
Work less than 20 hours per week for pay	5	8.47
Full or part-time student with no work for pay	1	1.69
Homemaker (unpaid)	6	10.17
Retired	3	5.08
Unemployed due to diagnosed emotional disability	10	16.95
Unemployed due to diagnosed physical disability	3	5.08
Unemployed for other reasons	2	3.39
Missing	2	3.39

21%). Co-therapy with both a mood stabilizer and an antidepressant was similar in both non-rapid-cycling patients and their rapid-cycling counterparts (41% vs. 43%).

Suicidality Risk

When the bipolar disorder patients who completed all the study surveys were assessed using the health system suicidality risk guideline, 19% of the respondents were evaluated as high risk, while 47% were considered moderate risk.

Work and Social Adjustment Scale

Almost 40% of the subjects worked at least 40 hours per week for pay (Table 2), while 17% reported being unemployed due to their emotional disability. A substantial proportion of bipolar disorder patients reported problems with work and social adjustment. While 42% reported that their work is at least markedly affected by their symptoms, over 50% reported experiencing difficulties with home management. Social activities were at least markedly affected in 37% of subjects, and 58% reported definite problems with private activities. Notably, about one quarter of these patients ranked problems with family and relationships as marked or severe.

When these respondents were asked, "How often did depressive symptoms affect activities/work?" 12% of these patients said all of the time, 35% said most of the time, and 29% said some of the time. Manic symptoms seemed to be less impactful, as only 14% of patients felt that their manic symptoms affected their activities/work all or most of the time and 35% said some of the time.

Fifty percent of these bipolar disorder patients reported missing at least 1 week of work (paid work/nonpaid work activities) during the past month. Even though 42% of respondents reported missing no days from work/school in the past 30 days due to depressive symptoms, approximately 1 in 5 (21%) reported having missed 10 days or more. Similarly, over half (57%) of respondents reported missing no days from work/school in the past 30 days due to manic symptoms, but 17% missed from 2 to 10 days. Overall, our work finds that subjects reported missing a mean of 7 days in the past month (absenteeism alone).

Table 3. Self-Reported Job Performance of Bipolar Disorder Patients Relative to Other Workers in Their Position

Job Performance Question	N	%
How often in the past 30 days was your performance higher than most workers in your job?		
All of the time	4	6.78
Most of the time	10	16.95
Some of the time	12	20.34
A little of the time	8	13.56
None of the time	20	33.90
Missing	5	8.47
How often in the past 30 days was your performance lower than most workers in your job?		
All of the time	4	6.78
Most of the time	11	18.64
Some of the time	13	22.03
A little of the time	12	20.34
None of the time	13	22.03
Missing	6	10.17

When absenteeism was subdivided by depressive and manic symptoms, the cohort reported a mean of 6.6 days in the past month lost from work or school due to depressive symptoms versus 3.9 days for manic symptoms.

This absenteeism appears to have had other effects as well, since 41% of these patients reported fearing the loss of their current job due to their emotional state, while 20% of subjects report being fired/laid off during the past 5 years due to their emotional state. Interestingly, when asked about higher and lower performance on the job relative to other workers in their position (Table 3), 47% felt that their performance was sometimes to all of the time lower than that of their peers. Similarly, in the past 30 days, 22% of the cohort reported not working as carefully as they should all or most of the time, and 17% felt that the quality of their work in the past 30 days was lower than it should have been most or all of the time. Work performance seems to vary by depressive and manic symptoms, as 75% of patients reported being less productive and achieving less in quantity or quality of work when experiencing depressive symptoms (vs. 36% during manic symptoms), while 44% report being more productive while experiencing manic symptoms (vs. 7% during depressive symptoms; Table 4).

DISCUSSION

Others have reported substantial underdetection of bipolar disorder in screened samples of primary care patients,^{1,2} which has brought to light the number of people who are misdiagnosed or overlooked. Our current work has sought to highlight more directly some of the key clinical (medication use, rapid cycling, suicidality) and social (labor force participation and social impact) features among those found to have bipolar disorder, which have not been adequately addressed in previous studies.

Table 4. Self-Reported Effect of Depressive and Manic Symptoms on Job Performance

Effect on Job Performance	Depressive Symptoms		Manic Symptoms	
	N	%	N	%
I notice no change in my ability to perform at work (or school or home)	6	10.17	3	5.08
I am less productive and achieve less in either quantity or quality of work	44	74.58	21	35.59
I am more productive and achieve more in either quantity or quality of work	4	6.78	26	44.07
I have not experienced these symptoms	3	5.08	7	11.86
Missing	2	3.39	2	3.39

One issue that has been suggested in the literature is the risk of rapid cycling and suicidality among those taking antidepressants, particularly SSRIs. Almost 1 in 4 subjects in our study (24%) were thought to be rapid cycling at the time of their visit; the psychiatry clinician felt that in 5 (36%) of the 14 rapid-cycling patients their rapid cycling was related to antidepressant use (7 were taking antidepressants). These findings are consistent with reports from the Stanley Foundation Bipolar German cohort, in which 40% of "rapid cyclers" suffered a recurrent rapid-cycling course during a 2.5-year follow-up period.⁹ A Veterans Affairs study cohort found that 14% of bipolar patients had experienced rapid cycling in the 12 months prior to intake (and 33% during the follow-up period),¹⁰ while Schneck and colleagues¹¹ report a rapid-cycling rate of 20% in their cohort of 500 bipolar patients.

The move to more aggressive recognition and treatment of depression, coupled with the underdetection of bipolar disorder among those presenting with depression, is probably a major explanatory factor in the rise of antidepressant use among bipolar patients.¹² Given that most patients with bipolar disorder present for treatment during the depressive phase rather than during manic or hypomanic phases, there is an increased likelihood of a unipolar depression diagnosis and treatment with antidepressants. The overuse of antidepressants has been associated with increased risk of antidepressant-induced mania, as 30% to 70% of patients with bipolar disorder are treated with antidepressants alone.¹³ However, Altshuler and colleagues¹⁴ found that bipolar patients who respond to antidepressants and do not switch into an episode of mania in the first 6 weeks are not at risk for switching over the ensuing year.

There are trials supporting the efficacy of antidepressants in bipolar disorder, along with evidence that they may induce rapid cycling, switching, and mania.¹⁵ Antidepressant-induced mania is often mixed, and since increased suicide risk is a particular feature of mixed states, this may explain why suicidal ideation can emerge with antidepressant treatment.¹⁵ The sentiment that antidepressants are "probably overused" and mood stabilizers "underused" has been expressed after critical review of the

studies in the nosology of bipolar disorder and the effects of antidepressant agents.¹⁶ About 1 in 5 members of our cohort (22%) experienced a mixed bipolar episode at the time of the screening. Furthermore, 61% of the cohort was found to be taking an antidepressant, almost half of whom (42%) were taking SSRIs, while 69.5% were also on treatment with a mood stabilizer.

Patients with bipolar disorder have been reported to have a higher risk of suicide than patients with any other psychiatric or medical illness,¹⁷⁻¹⁹ with rates of suicidal ideation and attempts between 35% and 50%.^{17,20,21} The completed suicide rate has been reported to be approximately 20%.²² Shi and colleagues,²³ in their study of recognized and unrecognized bipolar disorder in 25,460 California Medicaid (Medi-Cal) patients, found that unrecognized bipolar patients were nearly 4 times more likely to attempt suicide and 50% more likely to be hospitalized than nonbipolar patients. We found that 19% of bipolar patients in our study were considered to be at high risk of suicide and 47% at moderate risk, meaning that two thirds of our cohort was at substantial risk for suicide. Estimates of lifetime risk of suicide in patients with bipolar disorder range from 8% to 20% or 10 to 20 times that of the U.S. general population²⁴⁻²⁹ and may account for between 9% and 60% of all deaths among those with bipolar disorder.³⁰ Decreased suicide attempts have been documented among users of mood stabilizers, especially lithium and valproex.³⁰

Labor Force Participation

In the current study, 17% of bipolar subjects reported that they were unemployed due to their "emotional disability," while another 8.5% were unemployed for other reasons. This is consistent with reports from other studies. Zwerling and colleagues,³¹ using the National Health Interview Survey Disability Supplement of 1994 to 1995, found that only 56% of persons with bipolar disorder participated in the workforce, and the likelihood of workforce participation among bipolar subjects was substantially lower than among those without these disorders (OR 0.60, 95% CI 0.48 to 0.75) in multivariate models controlling for major demographics, education, race, and marital status. This would also seem to support the findings of Abood et al.,³² which refuted the commonly held notion that bipolar disorder does not necessarily confer social disadvantage (and is overrepresented in higher socioeconomic strata). Kupfer and colleagues²⁰ reported that 64% of those in the Stanley Center Bipolar Disorder Registry population of 2839 patients were unemployed, which is about 4 times the rate that we found. The difference is probably attributable to differences in the populations studied: the Stanley Registry recruits from those actively in treatment and those active in support groups, while our study population consists of those currently in treatment and insured for health care services, which likely reflects positive employment status.

Kessler and colleagues,³³ citing findings from the National Comorbidity Survey, found that patients with bipolar I experienced a mean 8.1 impairment days in the past month (bipolar II 3.8 days), which accounts for both absenteeism and presenteeism at work and at home; our research finds that subjects reported missing a mean of 7 days in the past month (absenteeism alone). When absenteeism was subdivided by depressive and manic symptoms, the cohort on average reported 6.6 days in the past month lost due to depressive symptoms versus 3.9 days for manic symptoms.

Results from the National Depressive and Manic-Depressive Association surveys suggest that 88% of those with bipolar disorder felt that their illness affected their ability to perform on their job, with only 40% reporting that they were employed at the time of the survey.³⁴ The current study provides additional data about the impact of depressive and manic symptom features separately, which adds a dimension to the data published to date.

The current study and its value must be viewed in the context of its potential limitations. First, the cohort was obtained from a series of patients with appointments in a psychiatry clinic that is part of a large managed care organization. This suggests that the subjects in this study may not be representative of the broad spectrum of all patients since they have access to health care through a managed care organization. Thus, it is possible that the impairment in this group is less than among those who may be unable to attain access to a managed care organization. Similarly, the coordination of care and the ease of referral from medical practitioners within a managed care setting probably increase likelihood of detection. Our protocol was explained to the staff and clinicians, and, as such, they were not blinded to the objectives of the study or the MDQ results of the patients, which may have biased our results and shown more favorable screening characteristics for the MDQ. Our validity calculations for the MDQ are higher than those reported by Hirschfeld⁵ in his original work; however, it is possible that the MDQ identifies lifetime experience versus a clinical diagnosis, which reflects current active disease. This may have led to a higher number of MDQ-positive but clinical diagnosis-negative patients in our study. Finally, a number of the patients in our study had been seen in this clinic before, which also increases the likelihood of detection and chronicity of disease. Although we had relatively high participation rates, we do not have comparator data.

This research documents some of the clinical features and social and labor-force impact of bipolar disorder in a managed care population and adds several dimensions to data published to date. More research is needed to continue the pursuit of better detection and treatment of bipolar disorder, which is extremely costly in individual, economic, and public-health terms.

Disclosure of off-label usage: The authors have determined that, to the best of their knowledge, no investigational information about pharmaceutical agents that is outside U.S. Food and Drug Administration–approved labeling has been presented in this article.

REFERENCES

- Kupfer DJ. Commentary: the increasing medical burden in bipolar disorder. *JAMA* 2005;293:2528–2530
- Das AK, Olfson M, Gameroff MJ, et al. Screening for bipolar disorder in a primary care practice. *JAMA* 2005;293:956–963
- Hirschfeld RMA, Cass AR, Holt DCL, et al. Screening for bipolar disorder in patients treated for depression in a family medicine clinic. *J Am Board Fam Pract* 2005;18:233–239
- Begley CE, Annegers JF, Swann AC, et al. The lifetime cost of bipolar disorder in the US: an estimate for new cases in 1998. *Pharmacoeconomics* 2001;19:483–495
- Hirschfeld RMA, Williams JBW, Spitzer RL, et al. Development and validation of a screening instrument for bipolar spectrum disorder: the Mood Disorder Questionnaire. *Am J Psychiatry* 2000;157:1873–1875
- Busch KA, Fawcett J, Jacobs DG. Clinical correlates of inpatient suicide. *J Clin Psychiatry* 2003;64:14–19
- Jacobs DG, Brewer M, Klein-Benheit M. Suicide assessment: an overview of recommended protocol. In: Jacobs DG, ed. *The Harvard Medical School Guide to Suicide Assessment and Intervention*. San Francisco, Calif: Jossey-Bass; 1998:3–39
- Mundt JC, Marks IM, Shear MK, et al. The Work and Social Adjustment Scale: a simple measure of impairment in functioning. *Br J Psychiatry* 2002;180:461–464
- Dittmann S, Biedermann NC, Grunze H, et al. The Stanley Foundation Bipolar Network: results of the naturalistic follow-up study after 2.5 years of follow-up in the German centres. *Neuropsychobiology* 2002;46(suppl 1):2–9
- Bauer MS, Kirk GF, Gavin C, et al. Determinants of functional outcome and healthcare costs in bipolar disorder: a high-intensity follow-up study. *J Affect Disord* 2001;65:231–241
- Schneck CD, Miklowitz DJ, Calabrese JR, et al. Phenomenology of rapid-cycling bipolar disorder: data from the first 500 participants in the Systematic Treatment Enhancement Program. *Am J Psychiatry* 2004;161:1902–1908
- Sachs GS. Bipolar mood disorder: practical strategies for acute and maintenance phase treatment. *J Clin Psychopharmacol* 1996;16(2 suppl 1):32S–47S
- Goodwin FK, Jamison KR. *Manic Depressive Illness*. New York, NY: Oxford University Press; 1990
- Altshuler L, Suppes T, Black D, et al. Impact of antidepressant discontinuation after acute bipolar depression remission on rates of depressive relapse at 1-year follow-up. *Am J Psychiatry* 2003;160:1252–1262
- Berk M, Dodd S. Are treatment emergent suicidality and decreased response to antidepressants in younger patients due to bipolar disorder being misdiagnosed as unipolar depression? *Med Hypotheses* 2005;65:39–43
- Ghaemi N, Sachs GS, Goodwin FK. What is to be done? controversies in the diagnosis and treatment of manic-depressive illness. *World J Biol Psychiatry* 2000;1:65–74
- Woods SW. The economic burden of bipolar disease. *J Clin Psychiatry* 2000;61(suppl 13):38–41
- Chen YW, Dilsaver SC. Lifetime rates of suicide attempts among subjects with bipolar and unipolar disorders relative to subjects with other axis I disorders. *Biol Psychiatry* 1996;39:896–899
- Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry* 1999;56:617–626
- Kupfer DJ, Frank E, Grochocinski VJ, et al. Demographic and clinical characteristics of individuals in a bipolar disorder case registry. *J Clin Psychiatry* 2002;63:120–125
- Suppes T, Leverich GS, Keck PE, et al. The Stanley Foundation Bipolar Treatment Outcome Network II: demographics and illness characteristics of the first 261 patients. *J Affect Disord* 2001;67:45–59
- Goldberg JF, Harrow M. Consistency of remission and outcome in bipolar and unipolar mood disorders: a 10-year prospective follow-up. *J Affect Disord* 2004;81:123–131
- Shi L, Thiebaud P, McCombs JS. The impact of unrecognized bipolar disorders for patients treated for depression with antidepressants in the fee-for-services California Medicaid (Medi-Cal) program. *J Affect Disord* 2004;82:373–383
- Bostwick JM, Pankratz VS. Affective disorders and suicide risk. *Am J Psychiatry* 2000;157:1925–1932
- Sharma R, Markar HR. Mortality in affective disorder. *J Affect Disord* 1994;31:91–96
- Guze SB, Robins E. Suicide and primary affective disorders. *Br J Psychiatry* 1970;117:437–438
- Harris EC, Barraclough B. Suicide as an outcome for mental disorders: a meta-analysis. *Br J Psychiatry* 1997;170:205–228
- Brodersen A, Licht RW, Vestergaard P, et al. Sixteen-year mortality in patients with affective disorder commenced on lithium. *Br J Psychiatry* 2000;176:429–433
- Fagiolini A, Kupfer DJ, Rucci P. Suicide attempts and ideation in patients with bipolar I disorder. *J Clin Psychiatry* 2004;65:509–514
- Goodwin FK, Fireman B, Simon GE, et al. Suicide risk in bipolar disorder during treatment with lithium and divalproex. *JAMA* 2003;290:1467–1473
- Zwerling C, Whitten PS, Sprince NL, et al. Workforce participation by persons with disabilities: the National Health Interview Survey Disability Supplement, 1994 to 1995. *J Occup Environ Med* 2002;44:358–364
- Aboud Z, Sharkey A, Webb M, et al. Are patients with bipolar affective disorder socially disadvantaged? a comparison with a control group. *Bipolar Disord* 2002;4:243–248
- Kessler RC, Mickelson KD, Barber CB, et al. The association between chronic medical conditions and work impairment. In: Rossi AS, ed. *Caring and Doing for Others: Social Responsibility in the Domains of Family, Work, and Community*. Chicago, Ill: University of Chicago Press; 2001:403–426
- Hirschfeld RM, Lewis L, Vornik LA. Perceptions and impact of bipolar disorder: how far have we really come? results of the National Depressive and Manic-Depressive Association 2000 Survey of Individuals With Bipolar Disorder. *J Clin Psychiatry* 2003;64:161–174

For the CME Posttest for this article, see pages 81–82.



Participants may receive a maximum of 1 *AMA PRA Category 1 Credit(s)*[™] by reading each CME article and correctly answering at least 70% of the questions in the Posttest that follows:

Go to www.psychiatrist.com/cmehome to take this Posttest online and earn credit immediately.

Or

1. Read each question carefully and circle the answer on the Registration Form.
2. Type or print the registration information in the spaces provided and complete the evaluation.
3. Send the Registration Form to the address or fax number listed on the Registration Form.

All replies and results are confidential. Answer sheets, once graded, will not be returned. Unanswered questions will be considered incorrect and so scored. The CME Institute of Physicians Postgraduate Press, Inc., will keep only a record of participation, which indicates the completion of the activity and the designated number of *AMA PRA Category 1 Credit(s)*[™] that have been awarded. Correct answers to the Posttest will be made available to the participants of this activity upon request after the submission deadline.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc., is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.



Answer to Pretest: 1. a

Stang et al.

pp. 42-47

1. In a 2005 primary care study, of those patients who screened positive for bipolar disorder, about _____% sought professional help for their symptoms and about _____% reported receiving a diagnosis of bipolar disorder.
 - a. 98; 72
 - b. 72; 8
 - c. 48; 45
 - d. 10; 8
2. In the present study, the Mood Disorder Questionnaire identified _____% of those with bipolar disorder (sensitivity) and _____% of those without (specificity).
 - a. 64; 94
 - b. 70; 59
 - c. 91; 94
 - d. 91; 59
3. When the patients with bipolar disorder who completed all the study surveys were assessed using the health system suicidality risk guideline, _____% of the respondents were evaluated as high risk for suicide, while _____% were considered moderate risk.
 - a. 4; 19
 - b. 47; 19
 - c. 9; 47
 - d. 19; 47
4. About _____% of patients with bipolar disorder worked at least 40 hours per week for pay, while about _____% were unemployed due to emotional disability.
 - a. 39; 17
 - b. 17; 8
 - c. 56; 17
 - d. 39; 8
5. Social activities were at least markedly affected in _____% of subjects.
 - a. 3
 - b. 17
 - c. 37
 - d. 73



REGISTRATION FORM

Circle the one correct answer for each question.

1. a b c d 4. a b c d
 2. a b c d 5. a b c d
 3. a b c d

Print or type

Name _____

Last 4 digits of Social Security Number _____
(Required to issue CME credit)

Birth Date (mm, dd, yy) _____
(Required to issue CME credit)

Degree _____ Specialty _____

Affiliation _____

Address _____

City, State, Zip _____

Phone () _____

Fax () _____

E-mail _____

- Hospital Private Practice Resident Intern

Deadline for submission

For a credit certificate to be issued, please complete this Registration Form no later than February 28, 2009. Online submissions will receive credit certificates immediately. Faxed or mailed submissions will receive credit certificates within 6 to 8 weeks.

Keep a copy for your files

Retain a copy of your answers and compare them with the correct answers, which will be published after the submission deadline.

Payment

If you complete the test online, no payment is necessary. A \$10 payment must accompany this form. You may pay by check, money order, or credit card (Visa or MasterCard). Make check or money order payable to Physicians Postgraduate Press, Inc. If paying by credit card, please provide the information below.

Check one: Visa MasterCard

Card number _____

Expiration date _____

Your signature _____

Please evaluate the effectiveness of this CME activity by answering the following questions.

- Was the educational content relevant to the stated educational objectives? Yes No
- Did this activity provide information that is useful in your clinical practice? Yes No
- Was the format of this activity appropriate for the content being presented? Yes No
- Did the method of presentation hold your interest and make the material easy to understand? Yes No
- Achievement of educational objective:
 - Enabled me to recognize clinical features of bipolar disorder and the impact of the disorder on patients' work and social lives. Yes No
- Did this CME activity provide a balanced, scientifically rigorous presentation of therapeutic options related to the topic, without commercial bias? Yes No
- Does the information you received from this CME activity confirm the way you presently manage your patients? Yes No
- Does the information you received from this CME activity change the way you will manage your patients in the future? Yes No
- If you answered yes, what change(s) do you intend to make in your practice?

- Please offer comments and/or suggested topics for future CME activities.

- How much time did you spend completing this CME activity?

- What is your preferred format for CME activities? Circle one.
 - Print media (e.g., journals, supplements, and newsletters)
 - Internet text
 - Internet multimedia
 - Audio CD
 - Live meeting
- Are you a physician? Yes No

TEAR OUT AND SEND THIS PAGE, ALONG WITH YOUR PAYMENT, TO:
 CME INSTITUTE • PHYSICIANS POSTGRADUATE PRESS, INC. • P.O. BOX 752870 • MEMPHIS, TN 38175-2870
 IF YOU ARE PAYING BY CREDIT CARD, YOU MAY FAX THIS PAGE TO: CME INSTITUTE AT 901-751-3444
 QUESTIONS? CALL 1-800-489-1001 EXT. 8 • WWW.PSYCHIATRIST.COM