

Update on Partial Response in Depression

Michael E. Thase, MD

Full symptomatic remission is the optimal outcome for patients with major depression. Unfortunately, antidepressant efficacy is limited to partial response for a significant minority of patients. Incomplete remission of depressive symptoms is associated with increased risk of relapse, decreased functioning in work and social settings, and increased risk of eventual suicide. Factors that increase the likelihood of incomplete remission include chronicity, severe symptomatology, and comorbid illnesses. Strategies to manage incomplete remission include “watchful waiting” (ie, continuing the original medication for another 4 to 8 weeks to see if complete remission will develop), switching antidepressants, or adding a second, adjunctive treatment (ie, either beginning psychotherapy or a second medication to augment the original antidepressant). Augmentation strategies may well prove to be the preferred strategy for improving response if tolerability is not an issue. Although studies on predictive factors have not yielded definitive results, clinicians in practice often select adjunctive agents that target patients’ persistent symptoms.

(J Clin Psychiatry 2009;70[suppl 6]:04–09)

Most, if not all, contemporary guidelines for the treatment of depression have adopted remission as the optimal outcome of an acute episode of depression.^{1,2} This decision is well-justified because patients with incomplete remission have increased risk of relapse,³ increased chronicity of depressive episodes with shorter durations between episodes,³ and impairments in workplace performance and social function when compared with those who are fully remitted.⁴ They also have increased all-cause mortality⁵ and risk of suicide.⁶ Essentially, the further a patient is from remission after treatment of an acute episode, the

more likely that he or she will have a poorer longer-term outcome.

CLINICAL REMISSION

Clinical remission of a depressive episode represents a nearly complete relief of the signs and symptoms of the presenting episode.⁷ Ideally, the patient will have no more symptoms of depression than someone who has never been depressed and, in the most practical terms, the individual will feel back to his or her usual self. A corollary of clinical remission is that, in addition to having virtually complete symptom relief, patients should also be able to return to normal levels of social or functional capacity.

Current definitions of remission are at the clinical level; they are not pathophysiologic. Looking toward the future of psychiatric therapeutics, an optimal treatment might also be hoped for to normalize the pathophysiology of an illness. However, that level of understanding of depression has not yet been reached.

Rates of remission, response without remission, and nonresponse vary from study to study. More than 2,700 patients were treated with citalopram for an average of 10 weeks as part of the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study.⁸ Only about one-third of the patients achieved remission at the end of the acute phase of treatment. Another approximately 14% improved but still had too many symptoms to be considered remitted; this group could be said to be partially remitted or to be responders without remission. The remainder—approximately one-half of those who received an adequate course of therapy—had either minimal or no improvement as a result of the initial treatment trial and might be classified as resistant to an adequate therapeutic trial of citalopram.

From the Departments of Psychiatry, University of Pennsylvania School of Medicine, Philadelphia Veterans Affairs Medical Center, and Pittsburgh Medical Center, Philadelphia and Pittsburgh.

This article is derived from the planning teleconference series “Tackling Partial Remission to Depression Treatment,” which was held in March and April 2009 and supported by an educational grant from Bristol-Myers Squibb Company and Otsuka America Pharmaceutical, Inc.


Dr Thase is a consultant for AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Forest, GlaxoSmithKline, Janssen, MedAvante, Neuronetics, Novartis, Pfizer, Schering-Plough, Sepracor, Shire US, Supernus, Transcept, and Wyeth; currently is a member of the speakers’ bureaus for AstraZeneca, Bristol-Myers Squibb, Eli Lilly, and Wyeth; has equity holdings in MedAvante; receives royalties from American Psychiatric Publishing, Inc, Guilford Publications, Herald House, and W.W. Norton & Company; has provided expert testimony for Jones Day (Wyeth litigation), Philips Lyttle, LLP (GlaxoSmithKline litigation), and Pepper Hamilton, LLP (Eli Lilly litigation); has received research funding (during the past 3 years) from Eli Lilly, Forest, GlaxoSmithKline, Sepracor, and the National Institute of Mental Health; and his spouse is an employee of Advogent, which does business with Bristol-Myers Squibb and Wyeth.

Corresponding author: Michael E. Thase, MD, University of Pittsburgh School of Medicine, 3535 Market St, Suite 670, Philadelphia, PA 19104 (thase@mail.med.upenn.edu).

doi:10.4088/JCP.8133su1c.01

© Copyright 2009 Physicians Postgraduate Press, Inc.

Figure 1. STAR*D Level 2 Remission Rates: Anxious vs Nonanxious Major Depressive Disorder^a




^aData from Fava et al.¹¹

**P* < .05.

Abbreviations: HDRS = Hamilton Depression Rating Scale, STAR*D = Sequenced Treatment Alternatives to Relieve Depression.

Figure 2. Impairment in Work and Relationships Normalizes Only With Remission^a



^aData from Miller et al.⁴

^bSignificant differences (*P* ≤ .05) existed in SAS work composite scores between response vs nonresponse, remission vs nonresponse, and remission vs response.

Abbreviation: SAS-SR = Social Adjustment Scale-Self Report.

the comorbid condition(s) is the next step. For example, if an individual experiencing depression also has a substance use disorder, making sure that he or she abstains from substance use may be the necessary intervention in order to ensure a full response to the antidepressant treatment. Likewise, recognizing and correcting a subtle thyroid dysfunction can be all the intervention that is necessary to see an individual with depression move from an incomplete response to a full remission.

Augmentation Versus Switching


Assuming that a depressed individual's symptoms have been properly tracked and measured, the dosage and

duration of the initial antidepressant have been optimized, and all comorbidities have been identified and treated to the extent that is possible, but the patient is still stuck in incomplete treatment response, then a couple of next-step strategies can be considered. One option is to simply switch to a second antidepressant medication. Although it is possible that another antidepressant would produce a complete remission, many clinicians reserve switching antidepressants for patients who have either obtained little symptom relief from the first medication or had trouble tolerating the side effects of an adequate dose of the first antidepressant. Thus, for a patient who has obtained definite symptom relief with the first antidepressant agent and has good tolerability, augmentation strategies probably rank higher than switching to another antidepressant.

An augmentation approach will build on the partial success of the first treatment, and, by continuing that initial treatment, patients do not have to go through medication withdrawal or cross-titration. The disruptive effects of medication withdrawal and cross-titration should not be minimized. Even if a patient has only improved by as little as 30%, it is remarkable how much the patient may miss that improvement if it is lost during the process of a cross-titration schedule. Thus, one potential advantage of an adjunctive strategy is that, by avoiding discontinuation and cross-titration, a therapeutic benefit may be produced more quickly than would be achieved with switching.

A second potential benefit of adjunctive strategy is that the second medication may be selected to match a patient's specific persistent residual symptoms. For example, an adjunctive medication with sedative/hypnotic properties could be added for a patient with persistent insomnia, or an adjunct medication with alertness/wakefulness-enhancing properties could be added for a patient with persistent fatigue or hypersomnolence.

Figure 3. Benefit-to-Risk Comparison for the Olanzapine-Fluoxetine Combination in Depression



Data from Thase et al.¹⁹

* $P < .001$ vs fluoxetine and olanzapine.

Abbreviation: MADRS = Montgomery-Asberg Depression Rating Scale.

PHARMACOTHERAPEUTIC AUGMENTATION OPTIONS

A meta-analysis¹³ of randomized, placebo-controlled lithium augmentation studies confirmed that lithium is an effective adjunctive treatment. Results varied considerably across studies, however, and most of these studies added lithium to tricyclic antidepressant (TCA) therapy. Lithium is one of the best studied adjunctive treatments but is not widely used today for the treatment of depression, perhaps because there are easier-to-implement strategies that do not entail a recommendation for blood level monitoring. Lithium should not be overlooked, however, and may be particularly valuable for patients with depressive syndromes that might be viewed as falling within the “softer” end of the bipolar spectrum.

A meta-analysis¹⁴ of studies of the use of thyroid hormone as an adjunct to TCA therapy also found evidence of efficacy. Thyroid augmentation is not as well studied as lithium augmentation and, as with lithium, most of the evidence comes from studies with TCAs. Larger studies are needed, with newer antidepressants. Whether thyroid hormone augmentation has specific antidepressant efficacy for patients taking selective serotonin reuptake inhibitors (SSRIs) or serotonin-norepinephrine reuptake inhibitors (SNRIs) remains to be seen, but the results of the STAR*D study¹⁵ showed fairly strong, although nonsignificant, trends favoring thyroid hormone augmentation over lithium augmentation in contemporary practice. This result may be because thyroid hormone augmentation is easier to implement. With respect to targeting a particular patient group, thyroid augmentation should be thought of whenever a patient who has not responded to antidepressants

is found to have low thyroid function, including those who have elevated thyroid-stimulating hormone levels.


Studies of modafinil have shown modest therapeutic benefit when added to antidepressant therapy, and to date the efficacy appears to be largely limited to the reduction of fatigue and daytime sleepiness.¹⁶ The net result is a small shift in the likelihood that patients receiving modafinil will move from being minimally improved or having responded without remission into remission.

With respect to conventional psychostimulants, the efficacy of methylphenidate augmentation has not yet been established. One small study¹⁷ suggested a beneficial trend, although none of the drug versus placebo differences reached statistical significance. Methylphenidate and other psychostimulants have the potential for abuse and misuse; therefore, these agents should be prescribed very carefully in off-label use as antidepressant therapy.

Atypical Antipsychotic Augmentation

Atypical antipsychotics are increasingly being used as augmenting strategies in the treatment of depression. Many clinicians are convinced that atypical antipsychotics have therapeutic benefit and that their efficacy is not limited to treating subtle psychosis, unrecognized bipolarity, sleep improvement, or anxiety reduction. Aripiprazole, in combination with many antidepressants, and olanzapine, in combination with fluoxetine, have received US Food and Drug Administration (FDA) approval for use in treating depression. A third atypical antipsychotic, quetiapine, has sufficient evidence and is currently under FDA review for this same indication. Perhaps the most important question about these medications is not whether they work but how long they should be maintained. Another important

Figure 4. Benefit-to-Risk Comparison for Aripiprazole in Depression

Data from Berman et al.²⁰* $P < .001$.

Abbreviation: MADRS = Montgomery-Asberg Depression Rating Scale.

question is whether these are the most cost-effective strategies to use, particularly for partial responders.

The efficacy of the olanzapine-fluoxetine combination was striking in the first small proof-of-concept study,¹⁸ but even a striking effectiveness profile must be balanced against the fairly high risk of weight gain with this therapy (Figure 3).¹⁹ Weight gain and the associated risk of metabolic side effects, such as dyslipidemia and glucose intolerance, need to be considered when using this antidepressant treatment strategy, and careful monitoring should be provided subsequently to minimize the potential for these adverse outcomes.

Aripiprazole was the first atypical antipsychotic approved for adjunctive use in patients not fully responding to antidepressants and is efficacious compared with placebo (Figure 4).²⁰ Aripiprazole has a lower risk of metabolic side effects—at least during short-term treatment—than does the olanzapine-fluoxetine combination, but aripiprazole does have one particularly troublesome set of adverse effects, namely, akathisia or less severe restlessness.²⁰

Augmentation Options for Comorbid Anxiety

Almost half of patients with major depression have comorbid anxiety,¹² and the presence of anxious symptoms in depression reduces the likelihood that patients will respond fully to antidepressant therapy. In the STAR*D study,¹¹ augmentation with the anti-anxiety agent buspirone was compared to augmentation with bupropion, a medication not known to be particularly useful for patients with a lot of anxiety. Curiously, and perhaps counterintuitively, trends—albeit nonsignificant ones—favored bupropion over buspirone among the patients with anxious depression (see Figure 1).

SUMMARY

Incomplete remission is an unfortunate and common outcome during the acute treatment of major depressive disorder and has become broadly recognized as a suboptimal outcome. Patients with complex presentations are at greater risk of incomplete remission. An important change of practice habits in the clinical setting would be to monitor the ongoing level of the symptomatic status of patients to best determine whether their response can be characterized as a full remission or a response without remission. In 2009, the art of practice for patients with depression who are tolerating their initial antidepressant well but have incomplete remission is to augment with psychotherapy and/or pharmacotherapy and select any pharmacotherapeutic adjunct on an individual basis to get the best balance of clinical benefit and minimum risk. A range of adjunctive strategies are now used to treat residual symptoms.

Drug names: aripiprazole (Abilify), bupropion (Aplenzin, Wellbutrin, and others), buspirone (BuSpar and others), citalopram (Celexa and others), fluoxetine (Prozac and others), imipramine (Tofranil and others), lithium (Lithobid, Eskalith, and others), methylphenidate (Concerta, Ritalin, and others), modafinil (Provigil), olanzapine (Zyprexa), olanzapine/fluoxetine (Symbyax), quetiapine (Seroquel), sertraline (Zoloft and others).

Disclosure of off-label usage: The author has determined that, to the best of his knowledge, buspirone, lithium, methylphenidate, modafinil, and quetiapine are not approved by the US Food and Drug Administration for the treatment of depression.

REFERENCES

1. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder. 2nd edition. http://www.psychiatryonline.com/pracGuide/loadGuidelinePdf.aspx?file=MDD2e_05-15-06. Accessed May 20, 2009.

2. Reesal RT, Lam RW, for the CANMAT Depression Work Group. Clinical guidelines for the treatment of depressive disorders, II: principles of management. *Can J Psychiatry*. 2001;46(suppl 1):21s–28s.
3. Judd LL, Paulus MJ, Schettler PJ, et al. Does incomplete recovery from first lifetime major depressive episode herald a chronic course of illness? *Am J Psychiatry*. 2000;157(9):1501–1504.
4. Miller IW, Keitner GI, Schatzberg AF, et al. The treatment of chronic depression, III: psychosocial functioning before and after treatment with sertraline or imipramine. *J Clin Psychiatry*. 1998;59(11):608–619.
5. Murphy JM, Monson RR, Olivier DC, et al. Affective disorders and mortality: a general population study. *Arch Gen Psychiatry*. 1987;44(5):473–480.
6. Judd LL, Akiskal HS, Paulus MP. The role and clinical significance of subsyndromal depressive symptoms (SSD) in unipolar major depressive disorder. *J Affect Disord*. 1997;45(1–2):5–18.
7. Thase ME. Introduction: defining remission in patients treated with antidepressants. *J Clin Psychiatry*. 1999;60(suppl 22):3–6.
8. Trivedi MH, Rush AJ, Wisniewski SR, et al. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry*. 2006;163(1):28–40.
9. Thase ME. Evaluating antidepressant therapies: remission as the optimal outcome. *J Clin Psychiatry*. 2003;64(suppl 13):18–25.
10. Rush AJ, Trivedi JH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry*. 2006;163(11):1905–1917.
11. Fava M, Rush AJ, Alpert JE, et al. Difference in treatment outcome in patients with anxious versus nonanxious depression: a STAR*D report. *Am J Psychiatry*. 2008;165(3):342–351.
12. Judd LL, Akiskal HS, Maser JD, et al. Major depressive disorder: a prospective study of residual subthreshold depressive symptoms as predictor of rapid relapse. *J Affect Disord*. 1998;50(2–3):97–108.
13. Crossley NA, Bauer M. Acceleration and augmentation of antidepressants with lithium for depressive disorders: two meta-analyses of randomized, placebo-controlled trials. *J Clin Psychiatry*. 2007;68(6):935–940.
14. Aronson R, Offman HJ, Joffe RT, et al. Triiodothyronine augmentation in the treatment of refractory depression: a meta-analysis. *Arch Gen Psychiatry*. 1996;53(9):842–848.
15. Nierenberg AA, Fava M, Trivedi MH, et al. A comparison of lithium and T(3) augmentation following two failed medication treatments for depression: a STAR*D report. *Am J Psychiatry*. 2006;163(9):1519–1530.
16. Fava M, Thase ME, DeBattista C. A multicenter, placebo-controlled study of modafinil augmentation in partial responders to selective serotonin reuptake inhibitors with persistent fatigue and sleepiness. *J Clin Psychiatry*. 2005;66(1):85–93.
17. Patkar AA, Masand PS, Pae CU, et al. A randomized, double-blind, placebo-controlled trial of augmentation with an extended release formulation of methylphenidate in outpatients with treatment-resistant depression. *J Clin Psychopharmacol*. 2006;26(6):653–656.
18. Shelton RC, Tollefson GD, Tohen M, et al. A novel augmentation strategy for treating resistant major depression. *Am J Psychiatry*. 2001;158(1):131–134.
19. Thase ME, Corya SA, Osuntokun O, et al. A randomized, double-blind comparison of olanzapine/fluoxetine combination, olanzapine, and fluoxetine in treatment-resistant major depressive disorder. *J Clin Psychiatry*. 2007;68(2):224–236.
20. Berman RM, Marcus RN, Swanink R, et al. The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: a multicenter, randomized, double-blind, placebo-controlled study. *J Clin Psychiatry*. 2007;68(6):843–853.