

Cardiac Safety Parameters of Olanzapine: Comparison With Other Atypical and Typical Antipsychotics

Joerg Czekalla, M.D.; Sara Kollack-Walker, Ph.D.;
and Charles M. Beasley, Jr., M.D.

Alterations of electrocardiogram results and cases of sudden cardiac death have been reported since the beginning of neuroleptic treatment. In particular, a temporal association exists between some antipsychotics and prolongation of the heart rate–corrected QT interval (QTc), an event that may increase the risk for developing a potentially fatal ventricular tachycardia arrhythmia known as torsades de pointes if it significantly exceeds normal intraindividual and interindividual variation. Although the incidence of serious adverse cardiac events in response to antipsychotic medications is relatively low, any possibility for the occurrence of cardiotoxicity warrants continued study. The present article reviews important differences among antipsychotic drugs in the potential for, and occurrence of, serious adverse cardiac outcomes and suggests that olanzapine, as therapeutically administered to patients with schizophrenia and related psychoses, does not contribute significantly to a QTc prolongation that could result in potentially fatal ventricular arrhythmias.

(*J Clin Psychiatry* 2001;62[suppl 2]:35–40)

Alterations of electrocardiogram results and cases of sudden cardiac death have been reported since the beginning of neuroleptic treatment.^{1,2} In particular, a temporal association exists between some antipsychotics and prolongation of the heart rate–corrected QT interval (QTc), an event that, if it significantly exceeds normal intraindividual and interindividual variation, may increase the risk for developing a potentially fatal ventricular tachycardia arrhythmia known as torsades de pointes.^{3,4} Although the incidence of serious adverse cardiac events in response to antipsychotic medications is relatively low, any possibility for the occurrence of cardiotoxicity warrants continued study. The present article reviews published data illustrating important differences among antipsychotic drugs in the potential for, and occurrence of, serious adverse cardiac outcomes. These data suggest that olanzapine is an atypical antipsychotic that, as therapeutically administered to patients with schizophrenia and related psychoses, does not contribute significantly to a

QTc prolongation that could result in potentially fatal ventricular arrhythmias.

MYOCARDIAL ACTION POTENTIAL, QTc PROLONGATION, AND MALIGNANT ARRHYTHMIAS

Recording of the surface electrocardiogram (ECG) reflects various phases of the myocardial action potential (Figure 1). In particular, the QT interval corresponds largely to the plateau phase of the myocardial action potential and is measured in milliseconds (ms) from the start of the QRS wave to the end of the T wave. The broad T wave of the ECG recording results from the rapid repolarization occurring nonsimultaneously throughout the ventricles of the heart. Because QT length shortens with increasing heart rate, the QT interval must be corrected for heart rate (QTc) using a regression formula (e.g., Bazett's formula [$QTcB = QT/RR^{1/2}$]).⁵

The length of the QTc interval can be influenced by numerous factors. For example, the QTc interval tends to be longer in females than in males⁶ and increases in length as a person ages.⁷ Considerable variation in the QTc interval has also been observed as a function of the time of day and even between ECG leads during testing,⁸ a phenomenon known as QTc dispersion. Furthermore, some individuals are born with a prolonged QTc interval that is referred to as the hereditary long QT syndrome (LQTS), while others develop what has been called an acquired LQTS (reviewed in Tan et al.⁹). Of interest, a variety of drugs, implicated in the etiology of acquired LQTS, have been shown


From the Medical CNS Division, Lilly Germany, Bad Homburg, Germany (Dr. Czekalla); and Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Ind. (Drs. Kollack-Walker and Beasley).

Supported by an unrestricted grant from Eli Lilly and Company.

Appreciation is expressed to Suraja Roychowdhury, Ph.D., and Jeff Ramsey, B.S., for technical assistance with the article and Cornelia Hoene for literature search.

Reprint requests to: Joerg Czekalla, M.D., Medical CNS Division, Lilly Germany, Saalburgstrasse 153, D-61350 Bad Homburg, Germany.

Figure 1. Effects of Antipsychotic Drugs on Cardiac Cells Can Prolong the QT Interval and Result in the Ventricular Arrhythmia Torsades de Pointes



to prolong the QTc interval, including certain antiarrhythmic drugs, antihypertensive drugs, anti-infective agents, tricyclic and tetracyclic antidepressants, and antipsychotic drugs (reviewed in Tan et al.⁹).

Prolongation of the QTc interval can predispose some patients to develop a polymorphic ventricular tachycardia known as torsades de pointes^{9,10} (see Figure 1). Torsades de pointes (twisting of points) is a unique ventricular tachycardia arrhythmia in which the mean electrical axis of the QRS complex appears to twist around the isoelectric line.¹¹ Torsades de pointes may serve as a transitional state between ventricular tachycardia and ventricular fibrillation, the latter representing a condition that if untreated can result in cardiac arrest. Although there is no consensus as to the length of the QTc interval below which the risk of arrhythmia is minimal, some expert opinion has suggested that a QTc of 500 ms is the lower limit for substantial risk of developing torsades de pointes.^{6,8}

Although a number of theories have been proposed, the occurrence of early afterdepolarizations within the myocardium is thought to be the main mechanism involved in QTc prolongation and development of ventricular arrhythmias (reviewed in Tan et al.⁹). Early afterdepolarizations can be triggered by various agents or conditions that reduce the magnitude of outward repolarizing K⁺ currents or enhance inward depolarizing Na⁺ or Ca²⁺ currents, leading to a delay of cardiac repolarization and the initiation of tachyarrhythmia. Drugs that significantly prolong the QTc interval or induce torsades de pointes have been found in *in vitro* experiments to alter these ion channel conductances.¹²

Antipsychotic Drugs and Cardiac Dysfunction

Antipsychotic drugs have been categorized into 2 main groups based primarily on differences in the appearance of

extrapyramidal symptoms (EPS). The majority of typical antipsychotic drugs are older drugs that fall into 3 main classes based on chemical structure: (1) phenothiazine derivatives (thioridazine, chlorpromazine, fluphenazine), (2) diphenylbutylpiperidine derivative (pimozide), and (3) butyrophenones (haloperidol, droperidol). As a group, the typical antipsychotics have been shown to be effective in treating the positive symptoms of schizophrenia (i.e., hallucinations and delusions), although with a fairly high incidence of EPS. In contrast, the newer atypical antipsychotics are as effective as the typical agents in treating positive symptoms of schizophrenia, apparently more effective in treating negative symptomatology as well as anxious and depressive symptoms accompanying schizophrenia, and have a superior EPS profile.¹³ Clinical availability of the atypical antipsychotics began with clozapine, to which risperidone, olanzapine, sertindole, quetiapine, and most recently ziprasidone (which is currently under review in several countries) were added subsequently.

Some antipsychotics have been shown to prolong the QTc interval, and as a consequence, their role in the development of torsades de pointes and sudden death has been suggested. However, recent data are beginning to suggest that there may be important differences among antipsychotics in their potential to cause cardiotoxicity. The reported effects of typical and atypical antipsychotics on electrocardiac function including mean QTc change, threshold QTc changes associated with torsades de pointes, and other cardiac safety reports are provided in Tables 1 and 2, respectively.

Typical Antipsychotics

Drugs within each class of the typical antipsychotics have been temporally associated with prolongation of the QTc interval, although to varying degrees. The largest mean increase in the QTc interval was observed for thioridazine (35.6 ms)^{14,70} and droperidol (59 ms, intravenous [i.v.] application),¹⁷ while the lowest increase was reported for haloperidol (range, 4.0–7.1 ms),^{15,16,70} with intermediate values reported for chlorpromazine¹⁵ and pimozide.¹⁶ Although only limited data have been provided for threshold change in QTc associated with torsades de pointes, one report¹⁸ shows that approximately 2% of patients treated with i.v. haloperidol show a QTc interval greater than 500 ms. Furthermore, numerous reports have indicated that each typical antipsychotic has been temporally associated with one or more serious cardiac events, with the majority of such events, including torsades de pointes, being observed after administration of the phenothiazine derivatives thioridazine and chlorpromazine (see Table 1).

A major concern that has been raised with the administration of typical antipsychotics is the clinical observation of a relationship between drug dose and QTc interval prolongation. In a recently published analysis (logistic

Table 1. Cardiac Safety Parameters Reported With Typical Antipsychotic Drugs^a

Typical Antipsychotic	Mean QTc Change ^b	Threshold QTc Changes Associated With Torsades de Pointes (% patients with QTc > 500 ms)	Cardiac Safety Reports
Thioridazine	33–35.8 ms ^{14,70} (oral)	Not reported	Clinical ECG data available since 1966 ² Reports of torsades de pointes, ^{19,20} malignant arrhythmia, ²¹ and sudden death ^{1,22} Greater increase in QTc with higher dose ^{4,14} Overdose was reported with cardiotoxicity ²³
Chlorpromazine	14 ms ¹⁵	Not reported	Clinical ECG data available since 1966 ² Reports of torsades de pointes ^{24,25} Greater increase in QTc with higher dose ⁴ Overdose (phenothiazines) was reported with cardiotoxicity ²³
Fluphenazine	Not reported	Not reported	Clinical ECG data available since 1965 ²⁶ Reports of T-ST wave abnormalities, supraventricular extrasystoles and broad QRS, ²⁷ and sudden death ²⁸ Greater risk of QTc prolongation with higher dose ⁴
Pimozide	19 ms ¹⁶	Not reported	Clinical ECG data available since 1987 ¹⁶ Reports of T-U wave abnormalities, ¹⁵ ECG changes in Tourette's disorder, ¹⁶ and sudden cardiac death in young adults ²⁹ Greater risk of QTc prolongation with higher dose ⁴
Droperidol	59 ms ¹⁷ (iv)	Not reported	Overdose was reported with torsades de pointes ³⁰ Clinical ECG data available since 1978 ³¹ Greater increase in QTc with higher dose ^{4,17}
Haloperidol	4.0–7.1 ms ^{15,16,70} (oral)	2% of patients QTc > 500 ms ¹⁸ (iv)	Overdose was reported with conduction disturbances and torsades de pointes ^{32,33} Clinical ECG data available since 1987 ¹⁶ Reports of torsades de pointes ^{18,34,35,36} and heart failure ³⁷ Greater increase in QTc with higher dose ³² Overdose was reported with QTc prolongation, life-threatening ventricular arrhythmia, and torsades de pointes ^{38–42}

^aAbbreviations: ECG = electrocardiogram, iv = intravenous, QTc = corrected QT interval.

^bRange of maximum mean QTc changes reported in different studies.

Table 2. Cardiac Safety Parameters Reported With Atypical Antipsychotic Drugs^a

Atypical Antipsychotic	Mean QTc Change ^b	Threshold QTc Changes Associated With Torsades de Pointes (% patients with QTc > 500 ms)	Cardiac Safety Reports
Clozapine	Not reported	Not reported	Clinical ECG data available since 1985 ⁵⁰ Reports of cardiomyopathy, ⁵¹ atrial fibrillation, ⁵² and sudden cardiac death ⁵³ Overdose was reported with decreased heart rate variability ⁵⁴
Risperidone	4.4–10.0 ms ^{43,70}	Not reported	Clinical ECG data available since 1997 ⁵³ Reports of cardiac arrest and QRS and QTc prolongation ⁵⁵ Overdose was reported with adverse cardiac effects ⁵⁶
Sertindole	24–30 ms ^{44,70}	3.1%–4.0% of patients QTc > 500 ms ⁴⁸	Clinical ECG data available since 1997 ⁴⁸ Reports of QTc prolongation and 3.5% arrhythmia vs 1.3% in reference group ⁵⁷ Greater increase in QTc with higher dose ⁴⁸
Olanzapine	–4.9 to 6.8 ms ^{43,45,70}	None ⁶⁹	Overdose was reported with torsades de pointes ^{58–60} Clinical ECG data available since 1997 ⁶¹ Overdose was reported with sinus tachycardia ^{62–64}
Quetiapine	8–14.5 ms ^{46,70}	Not reported	Clinical ECG data available since 1997 ⁴⁶ Reports of ECG changes ^{15,65} Overdose was reported with prolonged QTc and tachycardia ^{66–68}
Ziprasidone (not yet approved in all countries)	4.7–20.6 ms ^{47,70}	0.06%–1.2% of patients QTc > 500 ms ^{49,70}	Clinical ECG investigations performed in comparison to marketed atypical antipsychotics ⁷⁰ Reports of QT prolongation have led to delays in approval by the FDA ²⁹


^aAbbreviations: ECG = electrocardiogram, FDA = U.S. Food and Drug Administration, QTc = corrected QT interval.

^bRange of maximum mean QTc changes reported in different studies.

regression) of ECGs obtained from 495 psychiatric patients, robust predictors of QTc lengthening (defined from a healthy reference group as > 456 ms) included not only treatment with typical antipsychotics but also dose of antipsychotic.⁴ Indeed, the majority of reported cases of seri-

ous cardiac events or sudden death occurred in patients prescribed more than 50 mg/day of either haloperidol or droperidol and after an overdose with drugs in each class (phenothiazine derivatives, diphenylbutylpiperidine derivatives, butyrophenones).

Figure 2. Mean Change in QTc From Baseline Following Olanzapine Treatment at All Doses (N = 955) or at the Highest Dose (N = 319) in Acutely Psychotic Patients^a




^aData from Czekalla et al.⁶⁹ Abbreviation: QTc = corrected QT interval.

Atypical Antipsychotics

Similar to the typical antipsychotics, QTc prolongation has been reported for all atypical antipsychotics studied to date. However, differences among these drugs have been reported for the mean QTc increase and the percentage of patients showing QTc values > 500 ms. The largest mean increase in the QTc interval (range, 24–30 ms) was observed for sertindole,^{44,70} whereas olanzapine^{43,45,70} was temporally associated with the smallest change (range, -4.9 to 6.8) (see Table 2). In fact, in a large clinical trial comparing treatment efficacies of haloperidol with those of olanzapine, changes in mean QTc among the olanzapine dose groups were all decreases (range, -0.58 to -3.47 ms), and none were statistically different from control (Figure 2).⁶⁹ A review of baseline QTc values and change during olanzapine treatment in this trial and 2 other schizophrenia trials suggested normal random variability rather than a consistent drug effect.⁶⁹ Similarly, in a large clinical trial⁴³ comparing treatment efficacies of risperidone and olanzapine, risperidone-treated patients demonstrated a significant increase in QTc interval prolongation (4.4 ms) in comparison to olanzapine-treated patients who again showed a decrease (-4.9 ms). Very recently, a randomized study⁷⁰ in psychotic patients (N = 183) compared the effects of atypical and typical antipsychotics on the QTc interval at their estimated peak concentrations (T_{max}) following oral administration. The largest mean QTc prolongation was reported with thioridazine (35.8 ms), the next largest increase was caused by ziprasidone (20.6 ms), followed sequentially by quetiapine (14.5 ms), risperidone (10.0 ms), olanzapine (6.4 ms), and haloperidol (4.7 ms).⁷⁰ Only thioridazine and ziprasidone increased QTc by 75 ms or more in at least one study patient.⁷⁰

An increased risk of developing malignant arrhythmias has also been associated with elevations in the QTc inter-

Figure 3. Distribution of All Corrected QT (QTc) Intervals at Baseline (N = 2700) and Maximum QTc Intervals With Olanzapine Treatment (N = 1342) in Acutely Psychotic Patients^a



^aData from Czekalla et al.⁶⁹ Note that approximately 0.1% of patients at baseline showed QTc intervals > 500 ms, while no patient in the olanzapine-treated group had a QTc interval > 500 ms.

val > 500 ms. Among the atypical antipsychotics, sertindole was associated with the greatest percentage of patients possessing a QTc interval > 500 ms (3.1%–4.0%).⁴⁸ A lower incidence (0.06%–1.2%) has been reported with treatment with ziprasidone.^{49,70} Of interest, a recent clinical trial analysis of the distribution of QTc values in acutely psychotic patients revealed no evidence for QTc intervals > 500 ms in schizophrenia spectrum patients treated with olanzapine (Figure 3).⁶⁹ Furthermore, in the treatment of acute mania, no significant differences were noted between olanzapine- and placebo-treated patients with respect to the incidence of treatment-emergent QTc interval prolongation.⁷¹ Most recently, comparative analyses of cardiac function following treatment with several atypical antipsychotics have revealed no pathologic effects of olanzapine in regard to heart rate variability or QTc intervals > 500 ms.⁷²

Sertindole, which had the highest mean QTc increase and the greatest percentage of patients with a QTc interval > 500 ms, has been linked to torsades de pointes after overdose.^{58–60} Olanzapine and risperidone were suggested to have a favorable overdose profile in comparison to the phenothiazines and butyrophenones.⁶⁴ Other reports of cardiac events temporally associated with an overdose of the atypical drugs include tachycardia, ECG changes, and a decrease in heart rate variability (see Table 2).

Other Variables Affecting Cardiotoxicity

Determining the role that antipsychotic drugs may play in clinical cardiotoxicity can be difficult in lieu of other factors that can increase the risk for cardiac dysfunction.

For example, the effect of a given drug on the QTc interval may be exacerbated by an interaction with other drugs or by the presence of other conditions that can increase QTc prolongation including severe bradycardia, electrolyte disturbances, and cardiac, nutritional, endocrine, and central nervous system disorders (reviewed in Tan et al.⁹). Furthermore, cardiac safety concerns may increase with the treatment of patients who possess an inherited form of LQTS (e.g., Jervell and Lange-Nielsen syndrome, Romano-Ward syndrome; reviewed in Tan et al.⁹). Interestingly, even psychiatric illness itself, through alterations in parasympathetic and sympathetic autonomic regulation of heart rate variability, can contribute to cardiac liability.⁷³

CONCLUSION

The precise etiology of sudden death in patients receiving antipsychotic medication is uncertain, but prolongation of the QTc interval and its association with cardiac arrhythmias like torsades de pointes is a possible cause. Reports of serious adverse cardiac events temporally associated with parenteral application, overdose, and higher dosages of phenothiazines (thioridazine, fluphenazine, chlorpromazine), pimozide, and butyrophenones (haloperidol, droperidol) have been frequently observed, although such events are typically fewer in number, and less severe, in patients treated with atypical agents. The available data do not suggest that olanzapine, as therapeutically administered to schizophrenia spectrum patients, contributes significantly to a QTc prolongation that could result in malignant arrhythmias.

Drug names: chlorpromazine (Thorazine and others), clozapine (Clozaril and others), haloperidol (Haldol and others), olanzapine (Zyprexa), pimozide (Orap), quetiapine (Seroquel), risperidone (Risperdal), thioridazine (Mellaril and others).

REFERENCES

- Hollister LE, Kosek JC. Sudden death during treatment with phenothiazine derivatives. *JAMA* 1965;192:1035-1038
- Huston JR, Bell GE. The effect of thioridazine hydrochloride and chlorpromazine on the electrocardiogram. *JAMA* 1966;198:134-138
- Warner JP, Barnes TR, Henry JA. Electrocardiographic changes in patients receiving neuroleptic medication. *Acta Psychiatr Scand* 1996;93:311-313
- Reilly JG, Ayis SA, Ferrier IN, et al. QTc-interval abnormalities and psychotropic drug therapy in psychiatric patients. *Lancet* 2000;355:1048-1052
- Bazett HC. An analysis of the time relations of electrocardiograms. *Heart* 1920;7:353-370
- Garson A Jr. How to measure the QT interval: what is normal? *Am J Cardiol* 1993;72:14B-16B
- Dekker JM, Schoute EG, Klootwijk P, et al. Association between QT interval and coronary disease in middle-aged and elderly men: the Zutphen study. *Circulation* 1994;90:779-785
- Morganroth J, Brozovich FV, McDonald JT, et al. Variability of the QT measurement in healthy men, with implications for selection of an abnormal QT value to predict drug toxicity and proarrhythmia. *Am J Cardiol* 1991;67:774-776
- Tan HL, Hou CJ, Lauer MR, et al. Electrophysiologic mechanisms of the long QT interval syndromes and torsades de pointes. *Ann Intern Med* 1995;122:701-714
- Crumb W, Cavero I. QT interval prolongation by non-cardiovascular drugs: issues and solutions for novel drug development. *Pharm Sci Technol Today* 1999;2:270-280
- Dessertenne F. Considerations sur l'electrocardiogramme de la fibrillation ventriculaire. *Arch Mal Coeur Vaiss* 1966;59:263-272
- Davis AS. The pre-clinical assessment of QT interval prolongation: a comparison of in vitro and in vivo methods. *Hum Exp Toxicol* 1998;17:677-680
- Worrel JA, Marken PA, Beckman SE, et al. Atypical antipsychotic agents: a critical review. *Am J Health Syst Pharm* 2000;57:238-255
- Hartigan-Go K, Bateman DN, Nyberg G, et al. Concentration-related pharmacodynamic effects of thioridazine and its metabolites in humans. *Clin Pharmacol Ther* 1996;60:543-553
- Eastham JH, Jeste DV. Treatment of schizophrenia and delusional disorder in the elderly. *Eur Arch Psychiatry Clin Neurosci* 1997;247:209-218
- Fulop G, Phillips RA, Shapiro AK, et al. ECG changes during haloperidol and pimozide treatment of Tourette's disorder. *Am J Psychiatry* 1987;144:673-675
- Lischke V, Behne M, Doelken P, et al. Droperidol causes a dose-dependent prolongation of the QT interval. *Anesth Analg* 1994;79:983-986
- Sharma ND, Rosman HS, Padhi ID, et al. Torsades de pointes associated with intravenous haloperidol in critically ill patients. *Am J Cardiol* 1998;81:238-240
- Paoloni P, Ciliberti D, Blasi N, et al. Iatrogenic torsade de pointes induced by thioridazine. *Minerva Cardioangiol* 1992;40:245-249
- Kemper AJ, Dunlap R, Pietro DA. Thioridazine-induced torsades de pointes: successful therapy with isoproterenol. *JAMA* 1983;249:2931-2934
- Donatini B, Le Blaye I, Krupp P. Transient cardiac pacing is insufficiently used to treat arrhythmia associated with thioridazine. *Cardiology* 1992;81:340-341
- Whyman A. Phenothiazine death: an unusual case report. *J Nerv Ment Dis* 1976;163:214-217
- Buckley NA, Whyte IM, Dawson AH. Cardiotoxicity more common in thioridazine overdose than with other neuroleptics. *J Toxicol Clin Toxicol* 1995;33:199-204
- Ng AS, Johan A, Chia BL, et al. Chlorpromazine induced T wave and QT alternans: a case report and review. *Singapore Med J* 1982;23:110-113
- Ochiai H, Kashiwagi M, Usui T, et al. Torsade de pointes with T wave alternans in a patient receiving moderate dose of chlorpromazine: report of a case [in Japanese]. *Kokyu To Junkan* 1990;38:819-822
- Hassert GL Jr, Peterson AE, Burke JC. Comparison of T-wave changes in the dog following intravenous administration of chlorpromazine and fluphenazine. *Nature* 1965;207:765-766
- Dencker SJ, Johansson R, Lundin L, et al. High doses of fluphenazine enanthate in schizophrenia. *Acta Psychiatr Scand* 1978;57:405-414
- Moore MT, Book MH. Sudden death in phenothiazine therapy: a clinicopathologic study of 12 cases. *Psychiatr Q* 1970;44:389-402
- Welch R, Chue P. Antipsychotic agents and QT changes. *J Psychiatry Neurosci* 2000;25:154-160
- Krahenbuhl S, Sauter B, Kupferschmidt H, et al. Case report: reversible QT prolongation with torsades de pointes in a patient with pimozide intoxication. *Am J Med Sci* 1995;309:315-316
- Stanley TH. Cardiovascular effects of droperidol during enflurane and enflurane-nitrous oxide anaesthesia in man. *Can Anaesth Soc J* 1978;25:26-29
- Lawrence KR, Nasraway SA. Conduction disturbances associated with administration of butyrophenone antipsychotics in the critically ill: a review of the literature. *Pharmacotherapy* 1997;17:531-537
- Michalets EL, Smith LK, Van Tassel ED. Torsade de pointes resulting from the addition of droperidol to an existing cytochrome P450 drug interaction. *Ann Pharmacother* 1998;32:761-765
- Metzger E, Friedman R. Prolongation of the corrected QT and torsades de pointes cardiac arrhythmia associated with intravenous haloperidol in the medically ill. *J Clin Psychopharmacol* 1993;13:128-132
- Jackson T, Ditmanson L, Phibbs B. Torsade de pointes and low-dose oral haloperidol. *Arch Intern Med* 1997;157:2013-2015
- Di Salvo TG, O'Gara PT. Torsade de pointes caused by high-dose intravenous haloperidol in cardiac patients. *Clin Cardiol* 1995;18:285-290
- Iwahashi K. Significantly higher plasma haloperidol level during co-treatment with carbamazepine may herald cardiac change. *Clin Neuropharmacol* 1996;19:267-270
- Henderson RA, Lane S, Henry JA. Life-threatening ventricular arrhythmia

- (torsade de pointes) after haloperidol overdose. *Hum Exp Toxicol* 1991;10:59–62
39. O'Brien JM, Rockwood RP, Suh KI. Haloperidol-induced torsade de pointes. *Ann Pharmacother* 1999;33:1046–1050
 40. Yoshida I, Sakaguchi Y, Matsuishi T, et al. Acute accidental overdosage of haloperidol in children. *Acta Paediatr* 1993;82:877–880
 41. Aunsholt NA. Prolonged Q-T interval and hypokalemia caused by haloperidol. *Acta Psychiatr Scand* 1989;79:411–412
 42. Perrault LP, Denault AY, Carrier M, et al. Torsades de pointes secondary to intravenous haloperidol after coronary bypass. *Can J Anaesth* 2000;47:251–254
 43. Tran PV, Hamilton SH, Kuntz AJ, et al. Double-blind comparison of olanzapine versus risperidone in the treatment of schizophrenia and other psychotic disorders. *J Clin Psychopharmacol* 1997;17:407–418
 44. Zimbroff DL, Kane JM, Tamminga CA, et al, for the Sertindole Study Group. Controlled, dose-response study of sertindole and haloperidol in the treatment of schizophrenia. *Am J Psychiatry* 1997;154:782–791
 45. Barnes TRE, McPhillips MA. Critical analysis and comparison of the side-effect and safety profiles of the new antipsychotics. *Br J Psychiatry* 1999;174:34–43
 46. Arvanitis LA, Miller BG, for the Seroquel Trial 13 Study Group. Multiple fixed doses of "Seroquel" (quetiapine) in patients with acute exacerbation of schizophrenia: a comparison with haloperidol and placebo. *Biol Psychiatry* 1997;42:233–246
 47. Keck P Jr, Buffenstein A, Ferguson J, et al. Ziprasidone 40 and 120 mg/day in the acute exacerbation of schizophrenia and schizoaffective disorder: a 4-week placebo-controlled trial. *Psychopharmacol* 1998;140:173–184
 48. Product Monograph: Serolect (Sertindole), revised. Copenhagen, Denmark: Lundbeck; January 1997:39–43
 49. Tandon R, Harrigan E, Zorn SH. Ziprasidone: a novel antipsychotic with unique pharmacology and therapeutic potential. *J Serotonin Res* 1997;4:159–177
 50. Beier J, Hastrup S. ECG changes developing during clozapine treatment. *Ugeskr Laeger* 1985;147:619–620
 51. Leo RJ, Kreeger JL, Kim KY. Cardiomyopathy associated with clozapine. *Ann Pharmacother* 1996;30:603–605
 52. Low RA, Fuller MA, Popli A. Clozapine induced atrial fibrillation. *J Clin Psychopharmacol* 1998;18:170
 53. Suttman I, Ditter S, Landgraf R, et al. Clozapine and sudden death. *Lancet* 2000;355:842–843
 54. Rechlin T, Claus D, Weis M. Heart rate variability in schizophrenic patients and changes of autonomic heart rate parameters under treatment with clozapine. *Biol Psychiatry* 1994;35:888–892
 55. Ravin DS, Levenson JW. Fatal cardiac event following initiation of risperidone therapy. *Ann Pharmacother* 1997;31:867–870
 56. Kopala LC, Day C, Dillman B, et al. A case of risperidone overdose in early schizophrenia: a review of potential complications. *J Psychiatry Neurosci* 1998;23:305–308
 57. Wehnert A, Hale A, Kasper S, et al. EPOS: increasing our understanding of the treatment of schizophrenia. *Int J Psychiatry* 1997;1:197–202
 58. Lee AM, Knoll JL, Suppes T. The atypical antipsychotic sertindole: a case series. *J Clin Psychiatry* 1997;58:410–416
 59. Barnett AA. Safety concerns over antipsychotic drug, sertindole [news brief]. *Lancet* 1996;348:256
 60. Hale AS. A review of the safety and tolerability of sertindole. *Int Clin Psychopharmacol* 1998;13:65–70
 61. Beasley CM Jr, Tollefson GD, Tran PV. Safety of olanzapine. *J Clin Psychiatry* 1997;58(suppl 10):13–17
 62. Fogel J, Diaz JE. Olanzapine overdose. *Ann Emerg Med* 1998;32:275–276
 63. Cohen LG, Fatalo A, Thompson BT, et al. Olanzapine overdose with serum concentrations. *Ann Emerg Med* 1999;34:275–278
 64. Capel MM, Colbridge MG, Henry JA. Overdose profiles of new antipsychotic agents. *Int J Neuropsychopharmacol* 2000;3:51–54
 65. Misra LK, Erpenbach JE, Hamlyn H, et al. Quetiapine: a new atypical antipsychotic. *S D J Med* 1998;51:189–193
 66. McManus DQ, Arvanitis LA, Kowalczyk BB, for the Seroquel Trial 48 Study Group. Quetiapine, a novel antipsychotic: experience in elderly patients with psychotic disorders. *J Clin Psychiatry* 1999;60:292–298
 67. Hustey FM. Acute quetiapine poisoning. *J Emerg Med* 1999;17:995–997
 68. Gajwani P, Pozuelo L, Tesar GE. QT interval prolongation associated with quetiapine (Seroquel) overdose. *Psychosomatics* 2000;41:63–65
 69. Czekalla J, Beasley C, Berg P, et al. Cardiac safety profile of olanzapine based on preclinical and clinical ECG data [abstract]. *Eur Neuropsychopharmacol* 1999;9(5, suppl):S293
 70. Psychopharmacological Drugs Advisory Committee (PDAC). FDA Background Document on Zeldox (PDAC0700.M01). Rockville, Md: Food and Drug Administration; July 19, 2000
 71. Tohen M, Sanger TM, McElroy SL, et al, for the Olanzapine HGEH Study Group. Olanzapine versus placebo in the treatment of acute mania. *Am J Psychiatry* 1999;156:702–709
 72. Agelink MW, Majewski T, Wurthmann C, et al. Effects of newer atypical antipsychotics on autonomic neurocardiac function: a comparison between amisulpride, olanzapine, sertindole, and clozapine. *J Clin Psychopharmacol*. In press
 73. Choy AM, Lang CC, Roden DM, et al. Abnormalities of the QT interval in primary autonomic failure. *Am Heart J* 1998;136:664–671