

Adult ADHD Pharmacotherapy

Several medications are used to treat adult attention-deficit/hyperactivity disorder (ADHD), some of which may be augmented with psychotherapy or with other medications to successfully treat complicated presentations of ADHD.

Stimulant Treatment

Mixed amphetamine salts extended release (XR), dexamethylphenidate XR, the prodrug lisdexamfetamine, and OROS methylphenidate are the only stimulants indicated for the treatment of adult ADHD. However, immediate-release (IR) formulations of mixed amphetamine salts and methylphenidate have been shown in studies to be effective.

Both formulations of mixed amphetamine salts^{1,2} and methylphenidate^{3,4} as well as lisdexamfetamine⁵ have significantly reduced symptoms of ADHD compared with placebo. Further, mixed amphetamine salts IR¹ and dexamethylphenidate XR⁶ have significantly improved patients' ADHD Rating Scale scores compared with placebo. A dose-dependent relationship was not specifically found for mixed amphetamine salts (up to 60 mg/day), dexamethylphenidate XR (up to 40 mg/day), or lisdexamfetamine (up to 70 mg/day).

Although stimulants have abuse liability warnings, concern about stimulant abuse has decreased as newer formulations with slow-release mechanisms have become available. One study⁷ showed that OROS methylphenidate had no detection or likeability in subjects tested every hour for 10 hours after ingestion.

Side effects of stimulants include dry mouth, insomnia, appetite suppression, headache, edginess, and increased blood pressure and pulse. Stimulants may also uncover vocal or motor tics.

Nonstimulant Treatment

Atomoxetine, a selective norepinephrine reuptake inhibitor, is also indicated for treating adult ADHD and has been shown to be significantly superior to placebo in reducing ADHD symptoms.⁸

Side effects of atomoxetine include dry mouth, insomnia, decreased appetite, nausea, sexual difficulty, dizziness, and increased blood pressure and heart rate.

Nonpharmacologic Therapy

Psychosocial therapy may be necessary in the comprehensive treatment of adults with ADHD. For patients who partially respond to medication, adjunctive cognitive-behavioral therapy has been shown to reduce symptoms of ADHD as well as produce significantly more responders than continued medication alone.⁹

ADHD and Comorbid Disorders

ADHD often co-occurs with sleep, mood, anxiety, and substance use disorders (SUD). However, clinicians should always screen patients with ADHD for these conditions, and, if a patient has 1 or more co-occurring disorders, it is recommended that the most impairing disorder should be treated first. There are currently no approved guidelines for treatment of adult ADHD and comorbid mental health disorders.

Treatment for ADHD and some comorbidities can be concurrent, such as augmentation with an α_2 -agonist for sleep disorders or an antidepressant for depression. Anxiety symptoms may decrease with the resolution of ADHD symptoms.

For patients who have a current SUD, it is generally prudent to wait to fully assess ADHD symptoms until the patient has been abstinent for 1 week to 1 month. Additionally, no definitive correlation between stimulant treatment and increased use of alcohol, cigarettes, or drugs of abuse in patients with ADHD has been found.¹⁰ It is generally recommended to bring both the ADHD and the SUD under the treatment umbrella.

Take-Home Points

Both stimulants and nonstimulants are effective in treating adult ADHD. Clinicians should individualize treatment for their patients with ADHD considering the response to medication and the presence of comorbid disorders.

Drug names: amphetamine/dextroamphetamine (Adderall, Adderall XR, and others), atomoxetine (Strattera), dexamethylphenidate (Focalin XR and others), lisdexamfetamine (Vyvanse), methylphenidate (Concerta, Ritalin, and others).

Disclosure of off-label usage: Dr. Adler has determined that, to the best of his knowledge, immediate-release amphetamine dextroamphetamine and immediate-release methylphenidate are not approved by the U.S. Food and Drug Administration for the treatment of adult ADHD.

REFERENCES

- Spencer T, Biederman J, Wilens T, et al. Efficacy of a mixed amphetamine salts compound in adults with attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2001;58:775-782
- Weisler RH, Biederman J, Spencer TJ, et al. Mixed amphetamine salts extended-release in the treatment of adult ADHD: a randomized, controlled trial. *CNS Spectr* 2006;11:625-639
- Spencer TJ, Biederman J, Wilens T. A large, double-blind, randomized clinical trial of methylphenidate in the treatment of adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2005;57:456-463
- Biederman J, Mick E, Surman C, et al. A randomized, placebo-controlled trial of OROS methylphenidate in adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2006;59:829-835. Correction 2007;61:1402
- Adler LA, Goodman D, Kollins S, et al, on behalf of the 303 Study Group. A double-blind, placebo-controlled study of the efficacy and safety of lisdexamfetamine dimesylate in adults with attention-deficit/hyperactivity disorder. *J Clin Psychiatry* 2008;69:1364-1373
- Spencer TJ, Adler LA, McGough JJ, et al. Efficacy and safety of dexamethylphenidate extended-release capsules in adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2007;61:1380-1387
- Spencer TJ, Biederman J, Ciccone PE, et al. PET study examining pharmacokinetics, detection and likeability, and dopamine transporter receptor occupancy of short- and long-acting oral methylphenidate. *Am J Psychiatry* 2006;163:387-395
- Michelson D, Adler L, Spencer T, et al. Atomoxetine in adults with ADHD: two randomized, placebo-controlled studies. *Biol Psychiatry* 2003;53:112-120
- Safren SA, Otto MW, Sprich S, et al. Cognitive-behavioral therapy for ADHD in medication-treated adults with continued symptoms. *Behav Res Ther* 2005;43:831-842
- Faraone SV, Biederman J, Wilens TE, et al. A naturalistic study of the effects of pharmacotherapy on substance use disorders among ADHD adults. *Psychol Med* 2007;37:1743-1752

This ACADEMIC HIGHLIGHTS UPDATE section of The Primary Care Companion to The Journal of Clinical Psychiatry is derived from the ACADEMIC HIGHLIGHTS "ADHD and Comorbid Disorders in Adults" (J Clin Psychiatry 2008;69:1328-1335). This report was prepared by the CME Institute of Physicians Postgraduate Press, Inc., and supported by an educational grant from Eli Lilly and Company.

The ACADEMIC HIGHLIGHTS was chaired by Lenard A. Adler, M.D., New York University School of Medicine and New York VA Harbor Healthcare System, New York.

Financial disclosure: Dr. Adler has received grant/research support from Abbott, Cortex, Bristol-Myers Squibb, Merck, Novartis, Pfizer, Shire, Eli Lilly, Ortho-McNeil/Janssen/Johnson & Johnson, New River, Cephalon, and the National Institute on Drug Abuse; is a member of the speakers bureaus for Eli Lilly and Shire; is a member of the advisory boards and is a consultant for Abbott, Cortex, Novartis, Pfizer, Shire, Eli Lilly, Ortho-McNeil/Janssen/Johnson & Johnson, New River, Cephalon, Merck, Organon, Sanofi-Aventis, and Psychogenics; and has received royalty payments from New York University for license of adult ADHD scales and training materials.

The opinions expressed herein are those of the faculty and do not necessarily reflect the views of the CME provider and publisher or the commercial supporter.

© Copyright 2008
Physicians Postgraduate Press, Inc.

For the CME Posttest for this
ACADEMIC HIGHLIGHTS UPDATE,
see pages 499-500.