

# Earn the maximum AMA PRA Category 1 Credit<sup>TM</sup> for this activity online.

To receive up to 2.5 AMA PRA Category 1 Credits, take the Posttest and complete the extended evaluation online at PSYCHIATRIST.COM, keyword: PCC5. You must complete the Posttest and the entire evaluation online to receive the maximum credit.

To receive up to 2.0 AMA PRA Category 1 Credits, take the Posttest and send the completed Registration Form to the address or fax number listed.

You must correctly answer at least 70% of the questions in the Posttest to be awarded credit. Unanswered questions will be considered incorrect and so scored. Answer sheets, once graded, will not be returned. All replies and results are confidential. The CME Institute of Physicians Postgraduate Press, Inc., will keep only a record of participation, which indicates that you completed the activity and the number of AMA PRA Category 1 Credits you have been awarded. Correct answers to the Posttest will be available upon request after the submission deadline.

# **Accreditation Statement**

The CME Institute of Physicians Postgraduate Press, Inc., is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.



Answers to Pretest: 1. b 2. a

#### Katerndahl

pp. 376-383

- **1.** In studies of patients presenting with chest pain, the presence of panic disorder was associated with:
  - a. Atypical angina only
  - b. Typical angina only
  - c. Both atypical and typical angina
  - d. Neither atypical nor typical angina
- 2. Compared to patients with ischemic heart disease, patients with chest pain but normal angiogram results differ in:
  - a. Levels of pain
  - b. Associated sweating and dyspnea
  - c. Coping strategies used
  - d. Psychosocial stress
- **3.** Panic disorder should be suspected among patients presenting with chest pain unless they have:
  - a. No identifiable organic cause for the chest pain
  - b. A significant arrhythmia
  - c. Asymptomatic mitral valve prolapse
  - d. Normal cardiac testing

- 4. Which of the following decision tools should be used to diagnose panic disorder among patients with chest pain?
  - a. Panic disorder criteria in the DSM
  - b. Predictive models of panic disorder
  - c. Heart rate index calculation during an episode of chest pain
  - d. Presence of at least 2 of the following characteristics: female sex, age under 50 years, atypical chest pain, high anxiety levels, and lack of coronary artery disease
- 5. Which of the following statements about the usefulness of drug trials in patients with chest pain and possible panic disorder is true?
  - a. Response to sublingual nitroglycerin is specific to angina
  - b. A decrease in panic attack frequency with high-potency benzodiazepines cannot be due to placebo effect
  - c. Sertraline does not affect levels of pain in patients with noncardiac chest pain
  - d. Propranolol can correct false positive electrocardiogram results seen in patients with mitral valve prolapse
- Academic Highlights

pp. 391–402

- 6. Mr. R has acute bipolar depression and suicidal thoughts. In selecting a treatment, which of the following would help reduce the risk of suicide?
  - a. Antidepressant alone
  - b. Lithium plus an antidepressant
  - c. Antidepressant plus an atypical antipsychotic
  - d. Lamotrigine plus an antidepressant
- 7. Ms. T was discharged from the hospital after having a manic episode while taking a mood stabilizer and an atypical antipsychotic. She has experienced substantial weight gain since her treatment began. Which of the following strategies would be *least* useful?
  - a. Reduce the dose of both medications
  - b. Switch to a different antipsychotic
  - c. Discontinue atypical antipsychotic treatment
  - d. Switch to a different mood stabilizer
- 8. Which of the following practices helps keep patients with bipolar disorder well?
  - Taking baseline and follow-up waist and fasting glucose measurements to assess adverse events
  - b. Aggressively treating residual manic or depressive symptoms
  - c. Referring patients for adjunctive psychotherapy
  - d. All of the above
- 9. Among other treatments, the following can offset adverse neuroplastic changes in the brain: exercise, selective serotonin reuptake inhibitors, cognitive-behavioral therapy, glucocorticoid antagonists, electroconvulsive therapy, and antidepressants.
  - a. True
  - b. False
- **10.** All of the following messages should be communicated to your patients who are on maintenance treatment for bipolar disorder *except*:
  - a. Participate in psychotherapy
  - b. Exercise regularly
  - c. Take medication as prescribed
  - d. Take medication for the rest of your life

# REGISTRATION FORM

# Circle the one correct answer for each question.

1.	а	b	с	d	6.	а	b	с	d
2.	а	b	с	d	7.	а	b	с	d
3.	а	b	с	d	8.	а	b	с	d
4.	а	b	с	d	9.	а	b		
5.	а	b	с	d	10.	а	b	с	d

#### Deadline for submission

For a credit certificate to be issued, please complete this Registration Form no later than October 31, 2011. Online submissions will receive credit certificates immediately. Faxed or mailed submissions will receive credit certificates within 6 to 8 weeks.

#### Payment

If you complete the posttest online, no payment is necessary. A \$10 payment must accompany this form. You may pay by check, money order, or credit card. Make check or money order payable to Physicians Postgraduate Press, Inc. If paying by credit card, please provide the information below.

Check one:  Vis	a 🛛 MasterCard	🖵 AmEx	Discover	
Card number				
Expiration date				
Your signature				

#### Print or type

Name
Last 4 digits of Social Security Number
Birth Date (mm/dd/yy)
Degree Specialty
Affiliation
Address
City, State, Zip
Phone ( )
Fax ( )
E-Mail(Your credit certificate will be sent to this e-mail address)
☐ Hospital  ☐ Private Practice  ☐ Resident  ☐ Intern

# Send this page, along with your payment, to:

CME Institute • Physicians Postgraduate Press, Inc. P.O. Box 752870 • Memphis, TN 38175-2870

If you are paying by credit card, you may fax this page to: CME Institute at 901-751-3444

Questions? Call 1-800-489-1001, ext. 8

Please evaluate the effectiveness of this CME activity.	Poor	Needs Improvement	Average	Above Average	Excellent
1. The method of presentation held my interest and made the material easy to understand.					
2. This activity provided a balanced, scientifically rigorous presentation of therapeutic options related to the topic, without commercial bias.					
3. The educational content was relevant to the stated educational objectives.					
4. This activity helped me to:					
A. Determine whether screening for panic disorder is indicated in patients presenting with chest pain.					
B. Implement an individualized treatment plan designed to alleviate symptoms and ensure compliance so that the patient with bipolar disorder may achieve full remission.					
5. This activity confirmed the way I already manage my patients.					
6. This activity provided practical suggestions I can use in my practice.					
7. This activity provided information that will help me change my practice.					
8. What changes do you intend to make in your practice as a result of participating in this activity?					
<ul> <li>9. I need to know more about (suggest future topics):</li></ul>					
<ul> <li>Print media (e.g., journals, supplements, and newsletters)</li> <li>Internet</li> <li>12. Are you a licensed physician?</li> <li>Yes</li> <li>No</li> </ul>	et text	L Internet mu	ultimedia		Audio