

Electroconvulsive Therapy Decreases Dopamine D₂ Receptor Binding in the Anterior Cingulate in Patients With Depression: A Controlled Study Using Positron Emission Tomography With Radioligand [¹¹C]FLB 457

Tomoyuki Saijo, MD, PhD; Akihiro Takano, MD, PhD; Tetsuya Suhara, MD, PhD; Ryosuke Arakawa, MD, PhD; Masaki Okumura, MD; Tetsuya Ichimiya, MD, PhD; Hiroshi Ito, MD, PhD; and Yoshiro Okubo, MD, PhD

Objective: Electroconvulsive therapy (ECT) has been confirmed as one of the most effective treatments in drug-resistant major depression. However, the mechanism of ECT is still poorly understood. Although several lines of studies have focused on its effect on dopamine neurotransmission, the effects of ECT on dopamine D₂ receptors in a living human brain have not been investigated. Using positron emission tomography (PET) scans with the radioligand [¹¹C]FLB 457, we aimed to evaluate the effect of ECT on extrastriatal D₂ receptor binding in medicated patients with major depressive disorder (MDD).

Method: Seven patients with a DSM-IV diagnosis of MDD underwent PET scans before and after a series of 6–7 treatments with bilateral ECT. Eleven healthy controls were scanned for comparison. All participants were scanned at the National Institute of Radiological Sciences, Chiba, Japan, between November 2000 and September 2005. The parametric images of [¹¹C]FLB 457 binding were generated on the basis of a simplified reference tissue model. Voxel-based methods were used to assess the effect of ECT on D₂ receptor binding.

Results: There were no significant differences in D₂ receptor binding between patients with MDD and controls. All 7 patients showed clinical improvements in response to ECT treatment ($P < .001$). Significant changes in D₂ receptor binding, a mean of 25.2% reduction, were found in the right rostral anterior cingulate (AC) following ECT ($P < .001$).

Conclusions: Electroconvulsive therapy decreased D₂ receptor binding in the rostral AC in MDD patients responding to ECT. Our finding suggests that one of the biologic mechanisms of ECT could be related to dopaminergic alteration in the rostral AC.

J Clin Psychiatry 2010;71(6):793–799

© Copyright 2009 Physicians Postgraduate Press, Inc.

Submitted: September 27, 2008; accepted January 30, 2009.

Online ahead of print: December 15, 2009 (doi:10.4088/JCP.08m04746blu).

Corresponding author: Tetsuya Suhara, MD, PhD, Department of Molecular Neuroimaging, Molecular Imaging Center, National Institute of Radiological Sciences, 4-9-1, Anagawa, Inage-ku, Chiba, 263-8555, Japan (suhara@nirs.go.jp).

Electroconvulsive therapy (ECT) has been confirmed as one of the most effective treatments in drug-resistant major depression.¹ Among several neurotransmitter systems, such as the serotonergic, noradrenergic, and GABAergic neurotransmitter systems, on which ECT has been reported to have an effect,^{2,3} it has also been suggested that the dramatic and early improvements of severe psychomotor retardation and appetite loss after ECT were related to the dopaminergic effect.⁴ In addition, ECT has been widely used, with documented efficacy, in the treatment of different mental disorders involving dopaminergic dysregulation, such as mania, schizophrenia, and catatonia, and it has also led to prolonged improvement in some patients with Parkinson's disease and effectively treated neuroleptic malignant syndrome.¹ Such clinical effects suggest that dopamine neurotransmission plays a crucial role in the mechanism of the action of ECT.

Several lines of studies have focused on the effect of ECT on dopamine neurotransmission. Both animal and human studies^{5,6} have suggested that repeated ECT enhances dopamine-mediated behaviors. Some animal studies^{7,8} have indicated that both acute and chronic electroconvulsive shock (ECS) at the head increases dopamine release in the frontal cortex and the striatum. A human study⁹ also demonstrated increases in the concentrations of homovanillic acid in cerebrospinal fluid after ECT.

Animal behavioral studies^{10,11} have demonstrated that ECS stimulates dopamine D₂ receptor functions. However, only a few animal studies have investigated D₂ receptor binding after ECS, and the results have not been consistent.^{12,13} In addition, the effects of ECT on D₂ receptors in humans have not been investigated.

Recently, a positron emission tomography (PET) study¹⁴ using [¹⁸F]setoperone revealed that chronic ECS decreases serotonin 5-HT₂ receptors in nonhuman primates. In a human PET study,¹⁵ decreased regional cerebral glucose metabolism was reported after ECT, but the effect on specific neurotransmitter systems has not been studied.

Common symptoms of depression, such as anhedonia and amotivation, can be viewed as impairment in the control of dopaminergic reward and motivational systems.¹⁶ Those systems are reportedly closely connected with the dopamine mesolimbic-cortical pathway functionally and anatomically.¹⁷

Table 1. Clinical Characteristics of Patients

| Patient | Age, y ^a | Sex | Drug | Dose, mg/d | Length of Treatment, wk | No. of Episodes | Duration of Illness, mo | Length of Current Episode, mo | Current Clinical Features | GAF Score at Pre-ECT | HDRS Score | |
|---------|---------------------|-----|--------------------------|------------|-------------------------|-----------------|-------------------------|-------------------------------|---------------------------|----------------------|----------------------|-----------------------|
| | | | | | | | | | | | Pre-ECT ^b | Post-ECT ^c |
| 1 | 32 | M | Paroxetine hydrochloride | 40 | 8 | 3 | 8 | 2 | Melancholia | 51 | 16 | 4 |
| 2 | 53 | M | Fluvoxamine | 125 | 8 | 3 | 56 | 2 | Melancholia | 40 | 17 | 3 |
| 3 | 36 | F | Fluvoxamine | 100 | 20 | 3 | 31 | 5 | Melancholia | 35 | 28 | 13 |
| 4 | 35 | F | Fluvoxamine | 150 | 16 | 2 | 11 | 4 | Melancholia | 55 | 12 | 5 |
| 5 | 52 | M | Paroxetine hydrochloride | 40 | 70 | 3 | 21 | 1 | Melancholia | 31 | 26 | 11 |
| 6 | 58 | M | Paroxetine hydrochloride | 40 | 31 | 1 | 9 | 9 | Melancholia | 35 | 21 | 14 |
| 7 | 33 | M | Paroxetine hydrochloride | 40 | 47 | 2 | 20 | 2 | Melancholia | 45 | 16 | 2 |

^aMean (SD) age was 42.7 (11.1) years. ^bMean (SD) pre-ECT HDRS score was 19.43 (5.83). ^cMean (SD) post-ECT HDRS score was 7.43 (5.06).

Abbreviations: ECT = electroconvulsive therapy; F = female; GAF = Global Assessment of Functioning; HDRS = 21-item Hamilton Depression Rating Scale, Japanese version; M = male.

The radioligand [¹¹C]FLB 457 is a well-known dopamine D₂/D₃ antagonist that has picomolar in vitro affinity to both D₂ and D₃ receptors and has proven useful for the quantification of extrastriatal D₂ receptors in the human brain.¹⁸ We have previously demonstrated that the measurements of cortical D₂ receptor binding with [¹¹C]FLB 457 are reproducible and reliable.¹⁹ In the present study, using [¹¹C]FLB 457, we measured D₂ receptors in patients with major depressive disorder (MDD) before and after ECT to clarify the effect of ECT on the dopaminergic role in MDD.

METHOD

Subjects

Seven inpatients (mean age = 42.7 years, SD = 11.1 years; 5 men and 2 women; all right-handed) with MDD who had consented to ECT for clinical reasons, such as poor medication response and patient's own preference, at Nippon Medical School, Tokyo, or at Asai Hospital, Togane, Japan, and 11 age-matched healthy volunteers (mean age = 40.5 years, SD = 11.3 years; all men; all right-handed) participated in this study (Table 1).

The 7 patients met *DSM-IV* criteria for MDD. Based on conventional unstructured interviews and medical histories, we excluded patients with psychiatric disorders other than MDD, such as schizophrenia, substance abuse, anxiety disorder, and personality disorder. All patients were being treated with selective serotonin reuptake inhibitors (SSRIs), and their daily doses remained unchanged throughout the study (Table 1). Their median duration of treatment before ECT was 20 weeks (range, 8–70 weeks). Their medications for anxiety and sleep disturbances included lorazepam (1.5 mg/d) and flunitrazepam (1–2 mg/d).

The severity of symptoms was assessed using the Japanese version of the 21-item Hamilton Depression Rating Scale (HDRS)²⁰ on the same day as each PET scan. The HDRS was evaluated by 3 trained psychiatrists. The ratings were reviewed after the interviews; disagreements were resolved by consensus, and the consensus ratings were then used in this study. Each patient's Global Assessment of

Functioning²¹ score was assessed by the treating physicians on the day before the patient's first ECT session.

The healthy volunteers were recruited from the surrounding community. Based on unstructured psychiatric screening interviews, they were free of current and past psychiatric or major medical disease and had no relatives with neuropsychiatric disorders.

This study was approved by the Ethics and Radiation Safety Committee of the National Institute of Radiological Sciences, Chiba, Japan; the Ethics Committee of Nippon Medical School, Tokyo, Japan; and the Ethics Committee of Asai Hospital, Togane, Japan. After we provided complete explanation of the study, written informed consent was obtained from all patients and healthy volunteers.

Electroconvulsive Therapy Procedure

The patients received a series of 6–7 bilateral ECT sessions, 2–3 per week. Electroconvulsive therapy was administered with a square-wave, brief-pulse, constant-current device (Thymatron DGX; Somatics Inc, Lake Bluff, Illinois). Before the stimulus, subjects received atropine (0.4 mg), propofol (2.0 mg/kg), and succinylcholine (0.75 mg/kg), with dose adjustment as needed. Standard bifrontotemporal placement of electrodes was used for the treatment. Stimulus intensity was determined as a function of age, initially, and was increased until an adequate electroencephalographic seizure activity of 25 seconds or longer was obtained. Treatment was terminated when patients were judged by the treating physicians to have shown adequate symptom response.

Positron Emission Tomography Procedure

Patients underwent baseline PET scans with [¹¹C]FLB 457 on the day before their first ECT session. Follow-up PET studies were performed on the day after the last ECT session. Healthy volunteers were scanned once for comparison. All PET scans were conducted at the National Institute of Radiological Sciences, Chiba, Japan, between November 2000 and September 2005. Patients and healthy volunteers were placed in a supine position with eyes closed, and head-fixation devices (Fixter Instruments, Stockholm, Sweden)

with thermoplastic attachments made to fit the individuals were used.

Radioligand [^{11}C]FLB 457 was synthesized by O-methylation of the corresponding precursors with [^{11}C]methyl iodide with high specific radioactivity, which was obtained by a reduction of [^{11}C]CO₂ with LiAlH₄ in an inert atmosphere with specially designed equipment.²² The radiochemical purities were higher than 95%.

After a transmission scan with a ^{68}Ge - ^{68}Ga source, a bolus of 81.4–248.0 MBq of [^{11}C]FLB 457 was injected into the antecubital vein with a 20-mL saline flush. Specific radioactivity of [^{11}C]FLB 457 was 44.5–393.6 GBq/ μmol at the time of injection. Dynamic PET data were acquired for 90 minutes using CTI-Siemens ECAT EXACT HR+ (CTI-Siemens, Knoxville, Tennessee) in 3D mode without arterial sampling.

T1-weighted magnetic resonance imaging (MRI), by Philips Intera, 1.5 Tesla (Philips Medical Systems, Best, The Netherlands), was acquired from all patients and healthy volunteers. The scan parameters were 1-mm-thick 3D T1 images with a transverse plane (TR/TE 21/9.2 ms, flip angle 30°, matrix 256 × 256 mm).

Data Analysis

All emission scans were reconstructed with a Hanning filter cut-off frequency of 0.4 (full width at half maximum [FWHM] = 7.5 mm).

The parametric images of [^{11}C]FLB 457 binding were generated on the basis of a 3-parameter simplified reference tissue model.^{23,24} This model gives parametric images of R_1 (relative delivery of radioligand normalized to the cerebellum) and binding potential ($\text{BP}_{\text{ND}} = f_{\text{ND}} B_{\text{avail}} / K_D$), in which f_{ND} is the free fraction in the nondisplaceable compartment, B_{avail} is the density of receptors available to bind radioligand in vivo, and K_D is the dissociation constant for the radioligand.²⁵ The cerebellum was used as the reference region because of its negligible D₂ receptor density for calculation.²⁶ The cerebellar tissue concentration of radioactivity was obtained from regions of interest for the cerebellum, which were manually delineated on the individual MRIs coregistered to the PET summated images for 90 minutes using SPM2 (Wellcome Department of Cognitive Neurology, London, United Kingdom).

Analysis of parametric images of BP_{ND} and R_1 was undertaken using SPM2 to investigate differences in the brain.²⁷ A ligand-specific template image²⁸ was used to define the stereotactic transformation parameters for the BP_{ND} and R_1 images of [^{11}C]FLB 457. Normalized BP_{ND} and R_1 images were smoothed with a Gaussian filter to 8-mm full-width half-maximum.

Group Comparisons and Correlation

Statistical analyses of the BP_{ND} and R_1 data were performed with SPM2 using voxel-based paired or unpaired t tests. The BP_{ND} and R_1 differences were estimated for the following 3 contrasts: (1) patients before and after ECT, (2) healthy volunteers versus patients before ECT, and

(3) healthy volunteers versus patients after ECT. For those analyses, the significant voxel level threshold was set at $P < .001$, uncorrected, with a minimal cluster size of > 100 voxels. The resulting t values were converted to z scores, with brain locations reported as x , y , and z coordinates in Montreal Neurological Institute space with approximate Brodmann areas identified by mathematical transformation of SPM2 coordinates into Talairach space.^{29,30}

For significant regions, relation to the clinical effect was investigated. The percent changes in BP_{ND} values in significant clusters after ECT and the correlation between changes in BP_{ND} values and those of HDRS scores were evaluated. The following procedures were performed in SPM2: (1) the detected significant cluster was extracted as a volume of interest (VOI) and (2) individual mean BP_{ND} values in the VOI were acquired by applying the detected significant cluster extracted as a VOI to each of the normalized BP_{ND} images. The Spearman correlation method was used to examine the relationship between changes in BP_{ND} variables and changes in HDRS scores. We also investigated individual mean BP_{ND} values for controls and R_1 values for patients before and after ECT in the above VOI by applying the detected significant cluster extracted as a VOI to each of the normalized BP_{ND} images of controls and the normalized R_1 images of patients, respectively.

Statistical Analysis of Clinical Data

The paired t test was used to statistically analyze the difference between HDRS scores 1 day before the first session and 1 day after the last session of ECT. Values of $P < .05$ were considered significant.

RESULTS

All patients showed clinical improvement in response to ECT (Table 1). The baseline HDRS score was reduced from a mean \pm SD of 19.43 ± 5.83 to 7.43 ± 5.06 (paired t test: $t_6 = 8.92$, $P < .001$).

A voxel-based paired t test showed significant reduction of BP_{ND} in patients after ECT compared with patients before ECT in the right anterior cingulate (AC) (area 24), which extended rostrally to mesial regions of area 32 and to some voxels in area 10 (Table 2 and Figure 1). In contrast, no changes in R_1 were seen in the patients between pre-ECT and post-ECT scans. Significant increases of BP_{ND} in patients after ECT compared with patients before ECT were not found.

Voxel-based unpaired t tests demonstrated no significant differences in BP_{ND} either for healthy volunteers versus patients before ECT or for healthy volunteers versus patients after ECT.


In the post-ECT patients, there was a mean \pm SD reduction in D₂ receptor binding of $25.2\% \pm 7.5\%$ in the VOI compared with the pre-ECT patients (Figure 2). No significant correlations were observed between changes in BP_{ND} values and changes in HDRS scores. The mean \pm SD individual BP_{ND} values for controls and for patients before and after ECT were 0.81 ± 0.26 , 0.92 ± 0.28 , and 0.70 ± 0.27 , respectively.

Table 2. Effect of ECT on [¹¹C]FLB 457

| Region | Brodmann Area | Cluster Level | | | Voxel Level | | | | | MNI ^a | | | Talairach ^b | | |
|--------|---------------|----------------------|----------------------|------------------------|--------------------------|--------------------------|----------|----------|------------------------|------------------|----------|----------|------------------------|----------|----------|
| | | <i>P</i> (corrected) | <i>K_E</i> | <i>P</i> (uncorrected) | <i>P</i> (FWE-corrected) | <i>P</i> (FDR-corrected) | <i>t</i> | <i>z</i> | <i>P</i> (uncorrected) | <i>x</i> | <i>y</i> | <i>z</i> | <i>x</i> | <i>y</i> | <i>z</i> |
| rAC | 10/24/32 | <.001 | 109 | <.001 | .042 | .067 | 20.5 | 4.92 | <.001 | 6 | 40 | -4 | 5 | 35 | -3 |
| | | | | | .133 | .077 | 16.85 | 4.69 | <.001 | 6 | 48 | 10 | 5 | 44 | 8 |
| | | | | | .308 | .078 | 14.62 | 4.51 | <.001 | 4 | 42 | 0 | 3 | 38 | 0 |
| | | | | | 1 | .085 | 9.8 | 3.99 | <.001 | 8 | 54 | 8 | 7 | 49 | 6 |

^aThe MNI column contains spatial coordinates in MNI canonical magnetic resonance imaging. ^bThe Talairach column contains spatial coordinates in the Talairach atlas.³⁰

Abbreviations: FDR = false detective rate, FWE = family-wise error, *K_E* = number of voxels in the cluster, MNI = Montreal Neurological Institute, rAC = right anterior cingulate.

Figure 1. Map of *t* Values Showing the Voxels With Significant Decrease in [¹¹C]FLB 457 Binding Potential After Electroconvulsive Therapy (ECT)^{a,b}

^a*P* < .001, uncorrected.

^bDetected areas exceed an uncorrected *P* value of .001 with 100 or more contiguous voxels. In the left panel, all clusters throughout the whole brain are demonstrated in the "glass brain" in SPM2. The red arrow indicates a significant cluster of 109 voxels in the right anterior cingulate. The right panel shows the *t* statistic for the effect of ECT superimposed on the SPM2 canonical single-subject T1 image. The color scale shows the *t* values.

The mean \pm SD individual *R*₁ values for patients before and after ECT were 0.81 ± 0.06 and 0.79 ± 0.04 , respectively. The paired *t* test revealed that there were no changes in *R*₁ values in the right AC cluster in patients between the pre-ECT and post-ECT scans (*t*₆ = 1.13, *P* = .302).

DISCUSSION


The major finding of this study was that ECT induced a mean \pm SD significant decrease of $25.2\% \pm 7.5\%$ in D₂ receptor binding in the right rostral (AC).

Our previous study³¹ indicated that extrastriatal [¹¹C]FLB 457 was not sensitive to endogenous dopamine. In healthy volunteers, interindividual variance in BP_{ND} has been suggested to be due mainly to variability in B_{max} rather than K_D.³² In clinical PET studies, BP_{ND} is thus commonly used as an index of receptor density.³³ It is therefore reasonable to assume that the decreased [¹¹C]FLB 457 binding observed in this study reflects a reduction of D₂ receptor density in the anterior cingulate of patients with MDD after ECT. Providing that there was no neuronal cell loss after ECT,³⁴ the reduction of receptor density levels after ECT could be attributable

to mechanisms adaptive to increased dopamine levels in the synaptic cleft.^{7,9} The detected area, the rostral region of the AC, receives dense dopamine innervation principally from the ventral tegmental area.³⁵ Moreover, in the medial prefrontal cortex, which would include part of the detected area, dopamine was reported to show slow clearance from the synaptic cleft.³⁶ Since the density of D₂ receptors has been reported to be sensitive to synaptic concentrations of dopamine, D₂ receptors could be down-regulated in response to higher dopamine availability in the rostral AC.³⁷ Chronic stimulation with D₂ receptor agonists can lead to down-regulation of D₂ receptors,³⁸ and repeated ECS is reported to sensitize D₂ receptors.^{10,11} Taken together, our finding of decreased D₂ receptor binding in the rostral AC could be interpreted as a down-regulation of D₂ receptors after the ECT-induced dopamine release and D₂ receptor stimulation in the same area.

There has been only 1 animal study on D₂ receptors in the AC after ECS.¹² However, it was an autoradiographic study using [³H]spiroperidol, demonstrating that ECS had no effect on D₂ receptor binding in the AC. The differences in the method of ECT administration, the species

Figure 2. [¹¹C]FLB 457 Binding Potential of Right Anterior Cingulate in 11 Controls and 7 Depressed Patients Before and After 1 Course of Electroconvulsive Therapy (ECT)^a



^aHorizontal lines indicate means.

differences, and the ligand selectivity could explain the difference in results.

The rostral AC, connecting reciprocally to both the dorsal AC and the subgenual AC,³⁹ is considered to act as a bridge between attention and emotion⁴⁰ and to consist of the limbic-cortical circuits that maintain homeostatic mood control against stress. The region has been postulated as being one of the distinct targets that different modes of treatment modulate, resulting in a variety of complementary chemical and molecular adaptations and homeostatic effects that establish a normal mood state.⁴¹ In addition, anhedonia and amotivation, which are frequent symptoms of depression, have been hypothesized to be associated with a dysfunction of the subgenual prefrontal cortex (part of which is the rostral AC) to maintain tonic dopaminergic-dependent, reward-related activity.⁴² According to these lines of thought, one of the most relevant factors of the mechanism of antidepressant action would be the amelioration of the function of this region to regulate mesolimbic-cortical dopamine neuron activity.

We found no significant correlation between the changes of BP_{ND} in the rostral AC and the changes in HDRS score. The small sample size and the lack of diversity of patients made it difficult to investigate the correlation between ECT effects and clinical improvements. In a related study, Marano et al⁴³ reported that ECT increased brain-derived neurotrophic factor during a course of ECT and that the change was accompanied by a significant decrease in HDRS score. This finding indicates that D₂ receptor binding change might have been most significant not after, but during, a course of ECT; a correlation might have been detected between HDRS score change and BP_{ND} change during a course of ECT. Furthermore, it might have been valuable to assess the motivation/anhedonia or psychomotor retardation scores, in addition to HDRS scores, as those factors are reported to have close relations to the dopaminergic effect. Further studies are needed to clarify this issue.

Our finding—decreased D₂ receptor binding in the AC after ECT—might point to the mechanism of ECT, not the

mechanism of antidepressant action. Electroconvulsive therapy has a distinct efficacy in the treatment of different mental disorders, such as mania and schizophrenia, and has also produced improvement in some patients with Parkinson's disease.¹ To clarify this question of whether our finding might point to the mechanism of ECT or the mechanism of antidepressant action, investigations of the effect of ECT on D₂ receptors in patients with different mental disorders will be required.

Significant differences in [¹¹C]FLB 457 binding between patients with depression and control subjects were not detected in this study. Results of neuroimaging studies of striatal D₂ receptor binding in MDD have been inconsistent. For instance, increased dopamine D₂ receptor binding or sensitivity^{16,44–46} has been found in depressive and/or suicidal patients. Meanwhile, no difference in D₂ receptor binding was found in 2 recent studies.^{47,48} There had been, however, only 1 *in vivo* study measuring extrastriatal D₂ receptor binding of patients with MDD, and it demonstrated no difference in [¹¹C]FLB 457 binding between 7 people with depression and 7 healthy controls.⁴⁹ Our result was consistent with this study. Nevertheless, both studies may have been affected by methodological limitations, such as small sample size, differences in sex distribution of each group, and differences in medications. Further study is needed to clarify this issue.

In our study, the additional analysis of BP_{ND} in the right AC of controls revealed relatively large interindividual variability of BP_{ND} values (Figure 2). This result is supported by a former study⁵⁰ that demonstrated a mean ± SD BP_{ND} value in the AC of healthy volunteers of 0.79 ± 0.2 (N = 10; 7 men and 3 women; age, 22–38 y). In this respect, our finding that D₂ binding did not differ between patients with MDD and controls despite the fact that all BP_{ND} values in the right AC of patients with MDD had decreased after ECT might be explained.

Several confounding factors need to be considered for the interpretation of the present results. Since our sample size was small, our data might not assume normality of distribution. Therefore, we conducted additional analysis using a nonparametric voxel-based approach (SnPM; Wellcome Department of Cognitive Neurology, London, United Kingdom) to confirm our result. Nonparametric voxel-based analysis also revealed significant reduction of BP_{ND} in the right AC in patients after ECT compared with patients before ECT (Montreal Neurological Institute coordinates: x = 4, y = 44, z = 4; pseudo *t* = 4.85; *P* = .008).

Although ECT might alter blood flow,⁵¹ we could not detect a change in R₁ values between patients before and after ECT, suggesting that altered blood flow was unlikely to be responsible for the decreases in BP_{ND}. Our method for quantification required the use of the cerebellum as a reference region for estimation of nonspecific binding and free ligand concentration. Functional or structural abnormality of the cerebellum in mood disorders might hamper its use as a reference.^{52,53} There is also a possibility that ECT could induce differences in tracer kinetics for the cerebellum. In

addition, a more recent study⁵⁴ with [¹¹C]FLB 457 in rodents has shown that it cannot be excluded that a small percentage of the activity in the cerebellum may represent specific binding to D₂ receptors. However, cerebellar time-activity curves normalized by injected radioactivity did not differ significantly between patients before and after ECT.

We found no main effect of ECT or ECT-by-time interaction using repeated-measures analysis of variance with Greenhouse-Geisser correction (ECT: $F_{1,12} = 0.03$, $P = .865$; ECT-by-time interaction: $F_{2,28,22,4} = 0.281$, $P = .784$). It should be noted that there was no significant difference between the patients before and after ECT with respect to the injected specific radioactivity and injected mass in [¹¹C]FLB 457 (radioactivity injected: $t_6 = -0.81$, $P = .449$; mass of [¹¹C]FLB 457 injected: $t_6 = 0.90$, $P = .402$). The mean \pm SD values for injected mass for patients before and after ECT were 0.43 ± 0.17 μ g and 0.49 ± 0.17 μ g, respectively. Increased receptor occupancy of unlabeled radioligand could lead to lower BP_{ND} values, constituting a possible source of error. In simulation studies, for D₂ receptor density of the AC, an injected mass of 0.5 μ g has been shown to result in a radioligand occupancy of 5%, while an injected mass of 1.0 μ g corresponded to 8% occupancy.⁵⁰ On the basis of this analysis, the reported difference of 0.43–0.49 should account for 1% or 2% difference in BP_{ND}.

Regarding the possible contribution of atropine, propofol, and succinylcholine to our findings, only the effects of atropine and propofol against D₂ receptors have so far been investigated. Atropine has been reported not to occupy striatal or nigral dopamine D₂ receptors.⁵⁵ Propofol has been reported not to interact strongly with D₂ receptors (K_{50} values, affinity constant, for propofol binding to D₂ receptors > 150 μ M).⁵⁶ Taken together, it is unlikely that these medications used with ECT contributed to decreased D₂ receptor binding in the AC.

In this study, all patients were established on SSRI (fluvoxamine or paroxetine) treatment and were defined as treatment nonresponders on the basis of the clinical judgment of the attending physicians. Among depressive patients receiving paroxetine (modal daily dose, 50 mg; range, 30–50 mg), an increase in D₂ receptor binding has been reported in the anterior cingulate gyrus in treatment responders but not in nonresponders.⁵⁷ Paroxetine has also been reported to improve social functioning in A1+ allelic patients with posttraumatic stress disorder to a greater extent than in A1– allelic patients with posttraumatic stress disorder.⁵⁸ In that study, A1+ allelic patients with lower D₂ receptor density of basal ganglia, relative to A1– allelic patients, were speculated to respond to paroxetine via up-regulation of D₂ receptors. Chronic treatment of depressed patients with fluvoxamine (daily dose, 300 mg) has been shown not to alter D₂ receptor binding in the basal ganglia.⁵⁹ For patients with obsessive-compulsive disorder, repeated administration of fluvoxamine (mean \pm SD daily dose, 233 ± 50 mg) has been reported to increase D₂ receptor binding in the basal ganglia.⁶⁰ The relationship between the effect of fluvoxamine and changes in D₂ receptor binding in the AC has not been

clarified. Taken together, although the effects of SSRIs cannot be ruled out completely, it is unlikely that SSRIs contributed to the reduction in D₂ receptor binding in the AC of treatment nonresponders in the present study.

In conclusion, using PET with [¹¹C]FLB 457, we measured D₂ receptors in patients with MDD before and after ECT and found that ECT decreased D₂ receptor binding significantly in the rostral region of the right AC. The present findings suggest that the down-regulation of brain D₂ receptors might be induced after ECT as a compensatory change secondary to the modulation of the dopamine system in the region. We provide evidence that one of the biologic mechanisms of ECT could be related to dopaminergic alteration in the rostral AC.

Drug names: atropine (Atropen and others), fluvoxamine (Luvox and others), lorazepam (Ativan and others), paroxetine (Paxil, Pexeva, and others), propofol (Diprivan and others), succinylcholine (Quelicin and Anectine).

Author affiliations: Department of Neuropsychiatry, Nippon Medical School, Tokyo (Drs Saijo, Arakawa, Okumura, Ichimiya, and Okubo); Department of Molecular Neuroimaging, Molecular Imaging Center, National Institute of Radiological Sciences, Chiba (Drs Saijo, Takano, Suhara, Arakawa, Okumura, and Ito); and Department of Psychiatry, Asai Hospital, Togane (Drs Saijo and Ichimiya), Japan.

Potential conflicts of interest: None reported.

Funding/support: This study was supported by a Health and Labor Sciences Research Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H19-KOKORO-004) from the Japanese Ministry of Health, Labor and Welfare. The sponsors of the study had no role in the study design, the collection, analysis, and interpretation of data, the writing of the report, or the decision to submit the paper for publication.

REFERENCES

1. American Psychiatric Association Committee on Electroconvulsive Therapy. *The Practice of Electroconvulsive Therapy: Recommendations for Treatment, Training, and Privileging*, Second Edition. Washington, DC: American Psychiatric Association; 2001.
2. Nutt DJ. Relationship of neurotransmitters to the symptoms of major depressive disorder. *J Clin Psychiatry*. 2008;69(suppl E1):4–7.
3. Sanacora G, Mason GF, Rothman DL, et al. Increased cortical GABA concentrations in depressed patients receiving ECT. *Am J Psychiatry*. 2003;160(3):577–579.
4. Browning SM, Cowen PJ. Changes in mood, appetite and psychomotor retardation in depressed patients given ECT. *Br J Psychiatry*. 1986;149:371–373.
5. Cowen PJ, Nutt DJ, Green AR. Enhanced 5-hydroxytryptamine and dopamine-mediated behavioural responses following convulsions, pt 2: the effects of anaesthesia and current conditions on the appearance of enhanced responses following electroconvulsive shock. *Neuropharmacology*. 1980;19(9):901–906.
6. Berrios GE, Canagasabay AF. Depression, eye blink rate, psychomotor retardation, and electroconvulsive therapy—enhanced dopamine receptor sensitivity. *Convuls Ther*. 1990;6(3):224–230.
7. Glue P, Costello MJ, Pert A, et al. Regional neurotransmitter responses after acute and chronic electroconvulsive shock. *Psychopharmacology (Berl)*. 1990;100(1):60–65.
8. Yoshida K, Higuchi H, Kamata M, et al. Single and repeated electroconvulsive shocks activate dopaminergic and 5-hydroxytryptaminergic neurotransmission in the frontal cortex of rats. *Prog Neuropsychopharmacol Biol Psychiatry*. 1998;22(2):435–444.
9. Hoffmann G, Linkowski P, Kerkhofs M, et al. Effects of ECT on sleep and CSF biogenic amines in affective illness. *Psychiatry Res*. 1985;16(3):199–206.
10. Smith SE, Sharp T. Evidence that the enhancement of dopamine function by repeated electroconvulsive shock requires concomitant activation of D1-like and D2-like dopamine receptors. *Psychopharmacology (Berl)*. 1997;133(1):77–84.
11. Zarrindast MR, Sahebgharani M, Burnham WM. The effect of

- electroconvulsive shock seizures on behaviour induced by dopaminergic agonists and on immobility in the Porsolt test. *Eur Neuropsychopharmacol.* 2004;14(6):509–514.
12. Barkai AI, Durkin M, Nelson HD. Localized alterations of dopamine receptor binding in rat brain by repeated electroconvulsive shock: an autoradiographic study. *Brain Res.* 1990;529(1-2):208–213.
 13. Martin KF, Phillips I, Cheetham SC, et al. Dopamine D₂ receptors: a potential pharmacological target for nomifensine and tranlycypromine but not other antidepressant treatments. *Pharmacol Biochem Behav.* 1995;51(4):565–569.
 14. Strome EM, Clark CM, Zis AP, et al. Electroconvulsive shock decreases binding to 5-HT₂ receptors in nonhuman primates: an in vivo positron emission tomography study with [¹⁸F]setoperone. *Biol Psychiatry.* 2005;57(9):1004–1010.
 15. Nobler MS, Oquendo MA, Kegeles LS, et al. Decreased regional brain metabolism after ECT. *Am J Psychiatry.* 2001;158(2):305–308.
 16. Verbeek WJ, Berk M, Paiker J, et al. The prolactin response to sulpiride in major depression: the role of the D₂ receptor in depression. *Eur Neuropsychopharmacol.* 2001;11(3):215–220.
 17. Tomkins DM, Sellers EM. Addiction and the brain: the role of neurotransmitters in the cause and treatment of drug dependence. *CMAJ.* 2001;164(6):817–821.
 18. Olsson H, Halldin C, Swahn CG, et al. Quantification of [¹¹C]FLB 457 binding to extrastriatal dopamine receptors in the human brain. *J Cereb Blood Flow Metab.* 1999;19(10):1164–1173.
 19. Sudo Y, Suhara T, Inoue M, et al. Reproducibility of [¹¹C]FLB 457 binding in extrastriatal regions. *Nucl Med Commun.* 2001;22(11):1215–1221.
 20. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry.* 1960;23:56–62.
 21. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition. Washington, DC: American Psychiatric Association; 1994:32.
 22. Halldin C, Farde L, Högberg T, et al. Carbon-11-FLB 457: a radioligand for extrastriatal D₂ dopamine receptors. *J Nucl Med.* 1995;36(7):1275–1281.
 23. Lammertsma AA, Hume SP. Simplified reference tissue model for PET receptor studies. *Neuroimage.* 1996;4(3 pt 1):153–158.
 24. Gunn RN, Lammertsma AA, Hume SP, et al. Parametric imaging of ligand-receptor binding in PET using a simplified reference region model. *Neuroimage.* 1997;6(4):279–287.
 25. Innis RB, Cunningham VJ, Delforge J, et al. Consensus nomenclature for in vivo imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab.* 2007;27(9):1533–1539.
 26. Suhara T, Sudo Y, Okauchi T, et al. Extrastriatal dopamine D₂ receptor density and affinity in the human brain measured by 3D PET. *Int J Neuropsychopharmacol.* 1999;2(2):73–82.
 27. Friston KJ, Holmes AP, Worsley KJ. Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapp.* 1995;2(4):189–210.
 28. Meyer JH, Gunn RN, Myers R, et al. Assessment of spatial normalization of PET ligand images using ligand-specific templates. *Neuroimage.* 1999;9(5):545–553.
 29. Lacadie CM, Fulbright RK, Rajeevan N, et al. More accurate Talairach coordinates for neuroimaging using non-linear registration. *Neuroimage.* 2008;42(2):717–725.
 30. Talairach JTP. *Co-Planar Stereotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*. New York, NY: Thieme Medical Publishers; 1988.
 31. Okauchi T, Suhara T, Maeda J, et al. Effect of endogenous dopamine on extrastriatal [¹¹C]FLB 457 binding measured by PET. *Synapse.* 2001;41(2):87–95.
 32. Farde L, Hall H, Pauli S, et al. Variability in D₂-dopamine receptor density and affinity: a PET study with [¹¹C]raclopride in man. *Synapse.* 1995;20(3):200–208.
 33. Laruelle M. Dopamine transmission in the schizophrenic brain. In: Hirsh S, Weinberger D, eds. *Schizophrenia: Part Two, Biological Aspects*. Oxford, United Kingdom: Blackwell Publishing; 2003:365–387.
 34. Devanand DP, Dwork AJ, Hutchinson ER, et al. Does ECT alter brain structure? *Am J Psychiatry.* 1994;151(7):957–970.
 35. Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to behaviour. *Brain.* 1995;118(pt 1):279–306.
 36. Waymunt HK, Schenk JO, Sorg BA. Characterization of extracellular dopamine clearance in the medial prefrontal cortex: role of monoamine uptake and monoamine oxidase inhibition. *J Neurosci.* 2001;21(1):35–44.
 37. Norman AB, Battaglia G, Creese I. Differential recovery rates of rat D₂ dopamine receptors as a function of aging and chronic reserpine treatment following irreversible modification: a key to receptor regulatory mechanisms. *J Neurosci.* 1987;7(5):1484–1491.
 38. Stanwood GD, Lucki I, McGonigle P. Differential regulation of dopamine D₂ and D₃ receptors by chronic drug treatments. *J Pharmacol Exp Ther.* 2000;295(3):1232–1240.
 39. Mega MS, Cummings JL, Salloway S, et al. The limbic system: an anatomical, phylogenetic, and clinical perspective. *J Neuropsychiatry Clin Neurosci.* 1997;9(3):315–330.
 40. Pizzagalli D, Pascual-Marqui RD, Nitschke JB, et al. Anterior cingulate activity as a predictor of degree of treatment response in major depression: evidence from brain electrical tomography analysis. *Am J Psychiatry.* 2001;158(3):405–415.
 41. Mayberg HS. Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment. *Br Med Bull.* 2003;65(1):193–207.
 42. Drevets WC. Prefrontal cortical-amygdalar metabolism in major depression. *Ann N Y Acad Sci.* 1999;877:614–637.
 43. Marano CM, Phatak P, Vemulapalli UR, et al. Increased plasma concentration of brain-derived neurotrophic factor with electroconvulsive therapy: a pilot study in patients with major depression. *J Clin Psychiatry.* 2007;68(4):512–517.
 44. D'Haenen HA, Bossuyt A. Dopamine D₂ receptors in depression measured with single photon emission computed tomography. *Biol Psychiatry.* 1994;35(2):128–132.
 45. Shah PJ, Ogilvie AD, Goodwin GM, et al. Clinical and psychometric correlates of dopamine D₂ binding in depression. *Psychol Med.* 1997;27(6):1247–1256.
 46. Klimek V, Schenck JE, Han H, et al. Dopaminergic abnormalities in amygdaloid nuclei in major depression: a postmortem study. *Biol Psychiatry.* 2002;52(7):740–748.
 47. Parsey RV, Oquendo MA, Zea-Ponce Y, et al. Dopamine D(2) receptor availability and amphetamine-induced dopamine release in unipolar depression. *Biol Psychiatry.* 2001;50(5):313–322.
 48. Kuroda Y, Motohashi N, Ito H, et al. Effects of repetitive transcranial magnetic stimulation on [¹¹C]raclopride binding and cognitive function in patients with depression. *J Affect Disord.* 2006;95(1-3):35–42.
 49. Montgomery AJ, Stokes P, Kitamura Y, et al. Extrastriatal D₂ and striatal D₂ receptors in depressive illness: Pilot PET studies using [¹¹C]FLB 457 and [¹¹C]raclopride. *J Affect Disord.* 2007;101(1-3):113–122.
 50. Olsson H, Halldin C, Farde L. Differentiation of extrastriatal dopamine D₂ receptor density and affinity in the human brain using PET. *Neuroimage.* 2004;22(2):794–803.
 51. Takano H, Motohashi N, Uema T, et al. Changes in regional cerebral blood flow during acute electroconvulsive therapy in patients with depression: positron emission tomographic study. *Br J Psychiatry.* 2007;190:63–68.
 52. Bench CJ, Friston KJ, Brown RG, et al. The anatomy of melancholia: focal abnormalities of cerebral blood flow in major depression. *Psychol Med.* 1992;22(3):607–615.
 53. Shah SA, Doraiswamy PM, Husain MM, et al. Posterior fossa abnormalities in major depression: a controlled magnetic resonance imaging study. *Acta Psychiatr Scand.* 1992;85(6):474–479.
 54. Ahmad R, Hirani E, Grasby PM, et al. Effect of reduction in endogenous dopamine on extrastriatal binding of [¹¹C]FLB 457 in rat brain: an ex vivo study. *Synapse.* 2006;59(3):162–172.
 55. Alcock SJ, Hemsley KM, Crocker AD. Atropine acts in the ventral striatum to reduce raclopride-induced catalepsy. *Eur J Pharmacol.* 2001;424(3):179–187.
 56. Appadu BL, Strange PG, Lambert DG. Does propofol interact with D₂ dopamine receptors? *Anesth Analg.* 1994;79(6):1191–1192.
 57. Larisch R, Klimke A, Vosberg H, et al. In vivo evidence for the involvement of dopamine-D₂ receptors in striatum and anterior cingulate gyrus in major depression. *Neuroimage.* 1997;5(4, pt 1):251–260.
 58. Lawford BR, Young RM, Noble EP, et al. D₂ dopamine receptor gene polymorphism: paroxetine and social functioning in posttraumatic stress disorder. *Eur Neuropsychopharmacol.* 2003;13(5):313–320.
 59. Moresco RM, Colombo C, Fazio F, et al. Effects of fluvoxamine treatment on the in vivo binding of [F-18]FESP in drug naive depressed patients: a PET study. *Neuroimage.* 2000;12(4):452–465.
 60. Moresco RM, Pietra L, Henin M, et al. Fluvoxamine treatment and D₂ receptors: a pet study on OCD drug-naive patients. *Neuropsychopharmacology.* 2007;32(1):197–205.