



Supplementary Material

Article Title: Antidepressant Nonadherence in Routine Clinical Settings Determined From Discarded Blood Samples

Author(s): Ashlee M. Roberson, BA; Victor M. Castro, MS; Andrew Cagan, BA; and Roy H. Perlis, MD, MSc

DOI Number: 10.4088/JCP.14m09612

List of Supplementary Material for the article

1. [eTable 1](#) Comparison of Individuals With Routine Blood Draws Within 90 Days of Antidepressant Prescription to Those Without

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary eTable 1. Comparison of individuals with routine blood draws within 90 days of antidepressant prescription to those without.

Feature	Prescribed study antidepressant in January 2014 (n = 1,762)		Antidepressant level assessed 14-90 days after prescription (n = 109)		Comparison	
	%		%		X ² 1	P value
Male sex	29		38		3.56	0.17
Public insurance	30		38		2.28	0.13
Psychiatric provider	18		18		0.00	1.00
Specialist (versus primary care)	55		55		0.00	1.00
MDD diagnosis*	12		17		1.16	0.28
Initial prescription	19		18		0.00	1.00
2+ refills on prescription	73		74		0.04	0.84
	Mean	SD	Mean	SD	Student t	P value
Age (years)	52.25	16.83	56.02	17.05	2.24	0.02
Charlson comorbidity index	4.30	4.29	7.12	5.71	5.06	< 0.001

*MDD, major depressive disorder