



Supplementary Material

Article Title: The Epidemiology of Antisocial Behavioral Syndromes in Adulthood: Results From the National Epidemiologic Survey on Alcohol and Related Conditions-III

Author(s): Risë B. Goldstein, PhD, MPH; S. Patricia Chou, PhD; Tulshi D. Saha, PhD; Sharon M. Smith, PhD; Jeeseun Jung, PhD; Haitao Zhang, PhD; Roger P. Pickering, MS; W. June Ruan, MA; Boji Huang, MD, PhD; and Bridget F. Grant, PhD, PhD

DOI Number: 10.4088/JCP.15m10358

List of Supplementary Material for the article

1. [eTable 1](#) Odds Ratios (95% Confidence Intervals) of DSM-5 Antisocial Personality Disorder versus Adulthood Antisocial Behavioral Syndrome and 12-Month Substance Use, Mood, Anxiety, and Posttraumatic Stress Disorders
2. [eTable 2](#) Odds Ratios (95% Confidence Intervals) of DSM-5 Antisocial Personality Disorder versus Adulthood Antisocial Behavioral Syndrome and Other Lifetime Psychiatric Disorders

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary eTable 1. Odds Ratios (95% Confidence Intervals) of DSM-5 Antisocial Personality Disorder versus Adulthood Antisocial Behavioral Syndrome and 12-Month Substance Use, Mood, Anxiety, and Posttraumatic Stress Disorders

Comorbid Psychiatric Disorder	Adjusted for Sociodemographic Characteristics ^a	Adjusted for Sociodemographic Characteristics and Other Psychiatric Disorders ^b
Any substance use disorder	1.4 (1.26-1.64)	1.2 (1.08-1.44)
Alcohol use disorder	1.4 (1.16-1.67)	1.1 (0.93-1.42)
Nicotine use disorder	1.5 (1.33-1.71)	1.3 (1.13-1.49)
Any drug use disorder	1.6 (1.34-1.91)	1.1 (0.90-1.36)
Any mood disorder	1.8 (1.51-2.09)	1.2 (0.97-1.38)
Major depressive disorder	1.3 (1.10-1.59)	0.9 (0.72-1.11)
Dysthymia/persistent depressive disorder	1.5 (1.15-2.04)	1.2 (0.87-1.60)
Bipolar I disorder	2.9 (2.19-3.81)	1.6 (1.23-2.18)
Any anxiety disorder	1.8 (1.53-2.05)	1.1 (0.96-1.34)
Panic disorder	1.9 (1.46-2.58)	1.2 (0.92-1.65)
Agoraphobia	2.4 (1.68-3.37)	1.4 (0.96-1.98)
Social phobia	1.8 (1.40-2.41)	1.1 (0.82-1.44)
Specific phobia	1.6 (1.30-2.00)	1.2 (0.94-1.47)
Generalized anxiety disorder	1.8 (1.44-2.14)	1.0 (0.81-1.26)
Posttraumatic stress disorder ^c	2.4 (1.98-2.86)	1.5 (1.24-1.91)

Significant ($p < 0.05$) odds ratios are shown in **boldface**.

^a Odds ratios are adjusted for sex, age, race/ethnicity, education, family income, marital status, urbanicity, and geographic region.

^b Odds ratios are adjusted for sex, age, race/ethnicity, education, family income, marital status, urbanicity, geographic region, and all other substance use and psychiatric disorders.

^c Posttraumatic stress disorder was narrowly defined to require ≥ 3 subcriteria each for Criteria D and E to be considered positive, in addition to duration of ≥ 1 month and impairment or distress.

Supplementary eTable 2. Odds Ratios (95% Confidence Intervals) of DSM-5 Antisocial Personality Disorder versus Adulthood Antisocial Behavioral Syndrome and Other Lifetime Psychiatric Disorders

Comorbid Psychiatric Disorder	Adjusted for Sociodemographic Characteristics ^a	Adjusted for Sociodemographic Characteristics and Other Psychiatric Disorders ^b
Any substance use disorder	1.7 (1.40-2.03)	1.4 (1.16-1.71)
Alcohol use disorder	1.6 (1.39-1.83)	1.2 (1.00-1.35)
Nicotine use disorder	1.7 (1.54-1.98)	1.4 (1.22-1.60)
Any drug use disorder	1.9 (1.68-2.21)	1.4 (1.17-1.58)
Any mood disorder	1.9 (1.60-2.24)	1.3 (1.06-1.51)
Major depressive disorder	1.4 (1.19-1.62)	1.0 (0.85-1.19)
Dysthymia/persistent depressive disorder	1.5 (1.27-1.88)	1.1 (0.88-1.38)
Bipolar I disorder	2.6 (2.03-3.45)	1.5 (1.17-2.01)
Any anxiety disorder	1.7 (1.46-1.98)	1.0 (0.89-1.24)
Panic disorder	1.7 (1.29-2.18)	1.0 (0.80-1.38)
Agoraphobia	2.0 (1.46-2.86)	1.2 (0.86-1.69)
Social phobia	1.7 (1.34-2.16)	1.0 (0.77-1.28)
Specific phobia	1.6 (1.32-1.98)	1.1 (0.91-1.42)
Generalized anxiety disorder	1.9 (1.53-2.28)	1.1 (0.90-1.36)
Posttraumatic stress disorder ^c	2.4 (2.03-2.86)	1.6 (1.31-1.91)
Borderline personality disorder	2.7 (2.33-3.15)	1.9 (1.58-2.21)
Schizotypal personality disorder	2.7 (2.23-3.25)	1.8 (1.43-2.16)

Significant ($p < 0.05$) odds ratios are shown in **boldface**.

^a Odds ratios are adjusted for sex, age, race/ethnicity, education, family income, marital status, urbanicity, and geographic region.

^b Odds ratios are adjusted for sex, age, race/ethnicity, education, family income, marital status, urbanicity, geographic region, and all other substance use and psychiatric disorders.

^c Posttraumatic stress disorder was narrowly defined to require ≥ 3 subcriteria each for Criteria D and E to be considered positive, in addition to duration of ≥ 1 month and impairment or distress.