

It is illegal to post this copyrighted PDF on any website.

# Circadian Pattern of Deaths Due to Suicide in Intoxicated Alcohol-Dependent Individuals

Subhajit Chakravorty, MD<sup>a,b,\*</sup>; Rachel V. Smith, MPH, PhD<sup>c</sup>; Michael L. Perlis, PhD<sup>b</sup>; Michael A. Grandner, PhD<sup>d</sup>; and Henry R. Kranzler, MD<sup>a,b</sup>

## ABSTRACT

**Objective:** Alcohol intoxication and dependence are risk factors for suicide, a leading cause of death in the United States. We examined the hours of peak and nadir in completed suicides over a 24-hour period among intoxicated, alcohol-dependent individuals. We also evaluated suicide-related factors associated with intoxication at different times of the day.

**Methods:** We analyzed cross-sectional data from the 2003–2010 National Violent Death Reporting System provided by 16 US states. In the primary database, the deceased individuals' alcohol-dependent status was classified as "yes" or "no or unknown." We restricted the analysis to alcohol-dependent individuals with alcohol level data available (N = 3,661). The primary outcome measure was the reported time of death. Secondary outcome measures were predisposing and injury-related factors. Individuals were classified on the basis of their blood alcohol level (BAL) as heavy drinking (BAL<sub>H</sub> [≥ 80 mg/dL]) or non-heavy drinking (BAL<sub>O</sub> [ $< 80$  mg/dL]). The time of injury was divided into 1-hour bins, which were used to compute the incidence of suicide over 24 hours. We also evaluated the association between clinical factors and BAL<sub>H</sub> for each of six 4-hour time periods beginning at 00:01 hours.

**Results:** The majority (73.4%) of individuals showed evidence of alcohol consumption prior to committing suicide. BAL<sub>H</sub> was observed in 60.7% of all individuals. Peak incidences in suicide were identified at 21:00 for BAL<sub>H</sub> and 12:00 for BAL<sub>O</sub>, with nadirs at 05:00 and 03:00 hours, respectively. In a multivariable analysis, between 20:01 and 00:00 hours, BAL<sub>H</sub> was associated with more risk and protective factors than BAL<sub>O</sub>.

**Conclusions:** Identifying critical times and associated risk factors for suicidal behavior may contribute to suicide prevention efforts in intoxicated alcohol-dependent individuals.

*J Clin Psychiatry* 2018;79(6):17m11800

**To cite:** Chakravorty S, Smith RV, Perlis ML, et al. Circadian pattern of deaths due to suicide in intoxicated alcohol-dependent individuals. *J Clin Psychiatry*. 2018;79(6):17m11800.

**To share:** <https://doi.org/10.4088/JCP.17m11800>

© Copyright 2018 Physicians Postgraduate Press, Inc.

<sup>a</sup>Corporal Michael J. Crescenzo VA Medical Center, Philadelphia, Pennsylvania

<sup>b</sup>Perelman School of Medicine of the University of Pennsylvania, Philadelphia, Pennsylvania

<sup>c</sup>University of Louisville School of Nursing, Louisville, Kentucky

<sup>d</sup>University of Arizona College of Medicine, Tucson, Arizona

\*Corresponding author: Subhajit Chakravorty, MD, Corporal Michael J. Crescenzo VA Medical Center, Perelman School of Medicine, MIRECC, 2nd Fl, Postal Code 116, 3900 Woodland Ave, Philadelphia, PA 19104 (subhajit.chakravorty@uphs.upenn.edu).

Suicide caused more than 42,000 US deaths in 2014, making it the tenth leading cause of mortality.<sup>1</sup> In contrast to an overall decrease in the US mortality rate, the incidence of suicide is increasing. In 2014, firearms were the most common means of committing suicide in men (55.4%) and the second most common in women (31%), after poisoning (34.1%).<sup>2</sup> Studies have identified multiple suicide risk factors, including being male,<sup>3</sup> older,<sup>2</sup> unmarried,<sup>4</sup> and either unemployed or a blue-collar worker<sup>3,5,6</sup>; having a history of suicidal ideation or suicide attempts,<sup>7,8</sup> a chronic medical illness,<sup>3,9,10</sup> access to firearms,<sup>11</sup> and a psychiatric disorder such as depression, bipolar disorder, or alcohol dependence<sup>3,7,10,12,13</sup>; and recent alcohol consumption.<sup>3</sup>

Approximately one-third of suicide victims show evidence of recent drinking,<sup>4,14</sup> and heavy drinking, in particular, is linked to an increased risk of suicide.<sup>13,15–18</sup> Heavy drinking has been defined as the consumption of  $\geq 5$  drinks per day on  $\geq 5$  days within the past 30 days.<sup>19</sup> This drinking pattern is common among individuals with alcohol dependence, whose suicide rate is 60–120 times that of the general population.<sup>20</sup> Other risk factors for suicide among alcohol-dependent (AD) individuals include psychosocial (marital and occupational) dysfunction and comorbid chronic medical and psychiatric disorders.<sup>21–23</sup>

Another predictor of suicidal behavior is the time of day. Community-based studies evaluating circadian patterns of suicide have shown peaks and troughs across a 24-hour period. The most consistent timing of the peak has been 09:00–11:00 hours.<sup>24–28</sup> Other reported peaks are 12:00–15:00,<sup>24,26</sup> 15:00–18:00,<sup>29</sup> and 18:00–21:00.<sup>28,30</sup> In contrast, the circadian nadir for suicide has consistently been during the early morning hours (0:00–06:00), particularly around 02:00–03:00.<sup>24–27,30,31</sup> Studies have also examined sex, age, and season of the year as potential moderators of these circadian patterns. In one study, the peak incidence for males was 08:01–12:00 hours, whereas for females it was 20:01–24:00 hours.<sup>32</sup> Two studies showed that individuals 65 years and older had a peak incidence of suicide at 08:00–11:00, with the peak for younger individuals being later in the day.<sup>26,32</sup> A seasonal effect has also been seen, with peaks between 09:00–10:00 during winter and 14:00–15:00 during summer.<sup>27</sup> However, there has been little research on the circadian patterns of suicides in AD individuals, including whether in this population any of the aforementioned risk factors are associated with drinking and suicidal risk at certain times of the day.

Welte and colleagues<sup>33</sup> evaluated the relation between alcohol consumption and a diurnal pattern in 806 individuals who committed suicide between 1972 and 1984 in Erie County, New York. They found that the number of suicides in the subgroup with evidence of recent alcohol consumption was highest at night (n = 47), intermediate in the evening (n = 38), and lowest in the

You are prohibited from making this PDF publicly available.

- This study sought to identify periods of increased vulnerability to suicide during the 24-hour period in alcohol-dependent individuals, a population at an elevated risk for suicide, in order to optimize suicide prevention efforts.
- More intoxicated persons, as compared to non-intoxicated persons, committed suicide during the time period 20:01–00:00 hours, and they had more associated risk and protective factors for intoxication.

afternoon ( $n = 27$ ). Although 60% of individuals had blood alcohol levels  $\geq 0.1\%$ , the authors did not state how many had a diagnosis of alcohol dependence or how the time periods were defined. In a case series of AD individuals who attempted suicide in India ( $N = 30$ ), Prasanna and colleagues<sup>34</sup> identified a pattern of suicide attempts across the day. The highest proportion of attempts occurred during the morning hours (36.7%), followed by nighttime (26.7%) and the afternoon (16.7%).

We found no published data on circadian patterns of suicide in intoxicated AD individuals, associations with suicide risk factors, or information on the means of committing suicide. Thus, we examined variation in the incidence of completed suicides across the 24-hour period in AD individuals, using data from the National Violent Death Reporting System (NVDRS) of the Centers for Disease Control and Prevention (CDC). The availability of blood alcohol level (BAL) data in these individuals allowed us to compare the circadian pattern of suicides in intoxicated and non-intoxicated individuals. Analyzing demographic, clinical, and ballistic information enabled us to evaluate both risk and protective factors in these individuals. We sought to identify periods of increased vulnerability to suicide to focus suicide prevention efforts on the peak periods of risk for this population.

## METHODS

### Dataset

We obtained detailed case information from the NVDRS to investigate deaths by suicide between 2003–2010.<sup>35</sup> The NVDRS dataset was originally compiled by the CDC from 18 participating US states, but has now been expanded to 32 states.<sup>35</sup> Michigan and Ohio were among these 18 states and commenced with data collection in 2010. However, because state-level reporting of data commenced in 2011,<sup>36</sup> we excluded the data from these 2 states from the analysis ( $n = 61$ ), leaving data from 16 states in the analysis.

NVDRS pools information on the same incident from 4 major sources to provide an accessible, anonymous database. The primary sources of data are death certificates and reports from coroners or medical examiners, law enforcement officials, and forensic laboratories. The information that is collected includes the circumstances related to the suicide (eg, time of injury, presence of depression, and major life stresses like relationship or financial problems).<sup>35</sup> The

University of Pennsylvania's Institutional Review Board approved the conduct of this archival analysis project (IRB #815690).

### Subjects

In the primary database, the deceased individuals' AD status was classified as "yes" or "no or unknown." We extracted data only for individuals for whom this variable was coded as "present" and for whom alcohol-level data were available ( $N = 3,722$ , which was 10.5% of the total sample of 35,332). After excluding the 61 observations from Ohio and Michigan, our final analytic sample comprised 3,661 individuals.

### Variables

- **Demographic variables** including age, sex, race/ethnicity (non-Hispanic white, black/African-American, Hispanic/Latino, Asian, and other), education, and marital, veteran, and homelessness status were extracted from case reports.
- **Time of fatal injury** was the reported time when the act of suicide occurred. It was rounded to the nearest integer such that any time within the first 59 minutes of a particular hour was categorized as occurring within that hour.<sup>36</sup> In addition, we created six 4-hour time bins from the hourly information.<sup>25,32</sup> The categories were 00:01–04:00, 04:01–08:00, 08:01–12:00, 12:01–16:00, 16:01–20:00, and 20:01–00:00.<sup>32</sup>
- **Blood alcohol level** obtained from the postmortem examination records was recorded in mg/dL. The threshold for heavy drinking was set at  $BAL \geq 80$  mg/dL, ie, the legal limit for driving in most states and used by the CDC in reporting NVDRS data and in previous studies.<sup>37–40</sup> Individuals were classified into 1 of 3 drinker categories based on their blood alcohol levels: (1) none:  $BAL_{neg}$  (0 mg/dL); (2) moderate:  $BAL_{mod}$  ( $> 0$  and  $< 80$  mg/dL), and (3) heavy:  $BAL_H$  ( $\geq 80$  mg/dL).<sup>37–39</sup>
- **Drug screen** findings were obtained from the postmortem reports.
- **Likely predisposing factors** included prior treatment for a mental health problem, mental health diagnoses, current depressed mood, previous suicide attempts, physical health problems, job problems, recent legal problems, and financial problems.
- **Suicide injury-related factors** included the season of the year and means used. The seasons of the year were winter (December–February), spring (March–May), summer (June–August), and fall (September–November). The means of suicide included a firearm, poisoning, hanging, strangulation, suffocation, and other. Firearm-related variables included the firearm type; caliber; whether the firearm was stolen, stored locked, or stored loaded; and the owner of the gun.

### Data Analysis

Fisher exact test was used to compare the frequency of suicides for each hour of injury between  $BAL_{neg}$  and  $BAL_{mod}$  groups. Because  $BAL_{neg}$  and  $BAL_{mod}$  groups did not differ

**It is illegal to post this copyrighted PDF on any website.**

**Table 1. Demographic, Predisposing, and Lethal Self-Injury–Related Characteristics of the Sample**

Characteristic	n	Overall (n=3,661), n (%)	BAL <sub>H</sub> (n=2,222), n (%)	BAL <sub>O</sub> (n=1,439), n (%)	<i>P</i> <sup>a</sup>
<b>Demographics</b>					
Age, mean (SD), y	3,661	42.8 (13.3)	42.3 (12.8)	43.5 (14.1)	<b>.01</b>
Male	3,661	3,086 (84.3)	1,893 (85.2)	1,193 (82.9)	<b>.07</b>
Race	3,661				<b>.005</b>
White		3,391 (92.6)	2,072 (93.3)	1,319 (91.7)	
Black		142 (3.9)	68 (3.1)	74 (5.1)	
Other <sup>b</sup>		128 (3.5)	82 (3.7)	46 (3.2)	
Hispanic	3,654	226 (6.2)	135 (6.1)	91 (6.3)	.78
Marital status	3,643				.52
Married		1,432 (39.1)	885 (40.0)	547 (38.3)	
Never married		1,130 (31.0)	672 (30.4)	458 (32.0)	
Other <sup>c</sup>		1,081 (29.7)	656 (29.6)	425 (29.7)	
Education	1,793 <sup>d</sup>				.89
Less than high school		397 (22.1)	238 (21.3)	159 (23.6)	
High school graduate		797 (44.5)	514 (46.0)	283 (41.9)	
More than high school		599 (33.4)	366 (32.7)	233 (34.5)	
Veteran	3,501	687 (19.6)	407 (19.1)	280 (20.5)	.34
Homeless	3,557	48 (1.4)	29 (1.4)	19 (1.4)	1.00
<b>Likely predisposing factors</b>					
Previous treatment for mental health problem	3,661	1,643 (44.9)	879 (40.0)	764 (53.1)	<b>&lt;.0001</b>
Type of first mental diagnosis	1,718 <sup>d</sup>				<b>.005</b>
Depression/dysthymia		1,396 (81.3)	613 (79.1)	783 (83.0)	
Bipolar disorder		172 (10.0)	95 (12.3)	77 (8.2)	
Schizophrenia		36 (2.1)	24 (3.1)	12 (1.3)	
Anxiety disorder		43 (2.5)	17 (2.2)	26 (2.8)	
Posttraumatic stress disorder		23 (1.3)	8 (1.0)	15 (1.6)	
ADD or hyperactivity disorder		13 (0.8)	4 (0.5)	9 (1.0)	
Obsessive-compulsive disorder		4 (0.2)	3 (0.4)	1 (0.1)	
Other <sup>e</sup>		31 (1.8)	11 (1.4)	20 (2.1)	
Current depressed mood	3,661	1,817 (49.6)	723 (50.2)	1,094 (49.2)	.55
Previous suicide attempt	3,661	929 (25.4)	534 (24.0)	395 (27.5)	<b>.02</b>
Physical health problem	3,661	552 (15.1)	313 (14.1)	239 (16.6)	<b>.04</b>
Job problem	3,661	710 (19.4)	447 (20.1)	263 (18.3)	.17
Recent criminal legal problem	3,661	580 (15.8)	330 (14.9)	250 (17.4)	<b>.05</b>
Financial problem	3,661	588 (16.1)	357 (16.1)	231 (16.1)	1.00
<b>Suicide injury-related factors</b>					
BAL, mean (SD), mg/dL	3,661	131.4 (117.7)	208.7 (85.6)	12.1 (21.6)	NA
Season of the year	3,661				.88
Summer (June–Aug)		1,026 (28.0)	636 (28.6)	390 (27.1)	
Spring (March–May)		906 (24.8)	540 (24.3)	366 (25.4)	
Fall (Sept–Nov)		868 (23.7)	525 (23.6)	343 (23.8)	
Winter (Dec–Feb)		861 (23.5)	521 (23.5)	340 (23.6)	
Weapon used	3,656				<b>&lt;.0001</b>
Firearm		2,136 (58.4)	1,409 (63.5)	727 (50.6)	
Hanging, strangulation, suffocation		771 (21.1)	464 (20.9)	307 (21.4)	
Poisoning		481 (13.2)	208 (9.4)	273 (19.0)	
Other <sup>f</sup>		268 (7.3)	139 (6.3)	129 (9.0)	
Positive for any drug other than alcohol <sup>g</sup>	2,027 <sup>d</sup>	1,537 (75.8)	845 (70.0)	692 (84.4)	<b>&lt;.0001</b>
Opiates	2,536	337 (13.3)	153 (10.0)	184 (18.2)	<b>&lt;.0001</b>
Cocaine	2,554	278 (10.9)	168 (10.9)	110 (10.8)	.95
Marijuana	1,899	188 (9.9)	121 (10.0)	67 (9.7)	.81
Amphetamines	2,424	90 (3.7)	43 (2.9)	47 (5.0)	<b>.01</b>
Other drugs	2,210	975 (44.1)	512 (39.4)	463 (50.9)	<b>&lt;.0001</b>
Antidepressants	2,221	485 (21.8)	249 (18.4)	236 (27.3)	<b>&lt;.0001</b>

<sup>a</sup>Boldface indicates significance at  $\alpha = .05$ .

<sup>b</sup>Includes, but is not limited to, American Indian, Asian/Pacific Islander.

<sup>c</sup>Includes widowed, divorced, married but separated, and single (not otherwise specified).

<sup>d</sup>Greater than 10% of responses are missing; therefore, this was excluded from univariable and multivariable analyses.

<sup>e</sup>Specific data for this category were not available.

<sup>f</sup>Includes, but is not limited to, non-powder gun, sharp instrument, blunt instrument, fall, explosive, drowning, fire or burns, motor vehicle (eg, buses, motorcycles), other transport vehicle (eg, trains, planes, boats).

<sup>g</sup>Percentage displayed is the percentage of individuals tested who had a positive result.

Abbreviations: ADD = attention-deficit disorder, BAL = blood alcohol level, BAL<sub>H</sub> = heavy drinking (based on BAL), BAL<sub>O</sub> = non-heavy drinking (based on BAL), NA = not applicable.

at any time point at a Bonferroni-adjusted significance level ( $P < .002$ ), the 2 groups were combined into the BAL<sub>O</sub> group, to which BAL<sub>H</sub> was compared in all subsequent analyses.

Descriptive statistics were calculated for the overall sample and stratified into BAL<sub>H</sub> and BAL<sub>O</sub> subgroups. Fisher exact tests, extended Cochran-Mantel-Haenszel statistics, and analysis of variance (ANOVA) were employed to compare the subgroups on descriptive variables. Qualitative differences in the temporal patterns of suicides were assessed on an hourly basis using plots for the total sample and for the subgroups, with differences assessed with Fisher exact tests. Odds ratios were used to estimate the probability of suicide occurring at each hour for individuals classified as BAL<sub>H</sub> compared to BAL<sub>O</sub>.


We also used sociodemographic variables, predisposing factors, and suicide injury-related factors (the reference group being BAL<sub>O</sub>) to evaluate risk and protective factors for alcohol intoxication (BAL<sub>H</sub>) for the six 4-hour bins of time of injury. Because bivariate analyses revealed multiple associations between characteristics of interest and BAL<sub>H</sub> for the 6 individual time bins, we used 6 separate multivariable logistic regression models to evaluate the correlates of BAL<sub>H</sub> for the time bins, using the factors identified as significant ( $P < .05$ ) as predictors in the bivariate analyses. We used modified stepwise regression analyses to determine the final model and the likelihood ratio test and Akaike Information Criterion to select the best model for each time bin. Cochran-Mantel-Haenszel statistics and ANOVAs were used to evaluate differences among the means across the categories of alcohol consumption or 4-hour circadian bin during which the suicide occurred. Data analysis was conducted using SAS 9.4 software.<sup>41</sup>

## RESULTS

As shown in Table 1, the sample's mean age was 43 years (SD = 13.3), and it was primarily white ( $n = 3,391$ , 93%) and male ( $n = 3,086$ , 84%). Twenty-five percent ( $n = 929$ ) of the sample had previously attempted suicide. The majority (73.4%) of individuals showed evidence of alcohol consumption prior to committing suicide. The average BAL at the time of death for the



**Figure 1. Proportion of Suicides That Occur per Hour by BAL at Death**



Abbreviations: BAL = blood alcohol level, BAL<sub>H</sub> = heavy drinking (based on BAL), BAL<sub>O</sub> = non-heavy drinking (based on BAL).

BAL<sub>H</sub> group was 208.7 mg/dL (SD = 85.6). The majority (60.7%) of suicide victims were in the BAL<sub>H</sub> category. More than half of the suicides ( $n = 2,136$ , 58%) were committed with a firearm. Among individuals tested for drug use following the suicide ( $n = 2,027$ , 55.4%), 76% tested positive ( $n = 1,537$ ) for at least 1 of the following drugs: amphetamines ( $n = 90$ , 4%), cocaine ( $n = 278$ , 11%), marijuana ( $n = 188$ , 10%), opiates ( $n = 337$ , 13%), or other drugs ( $n = 975$ , 44%). Twenty-two percent ( $n = 485$ ) of individuals screened positive for antidepressant use during postmortem examination. Notably, the BAL<sub>O</sub> group had a significantly higher proportion of individuals with a history of mental health treatment (53% vs 40%,  $P < .0001$ ), a previous suicide attempt (28% vs 24%,  $P = .02$ ), and a positive toxicologic test at suicide (84% vs 70%,  $P < .0001$ ).

As shown in Figure 1, overall, the highest proportion of suicides occurred in the latter half of the day (ie, between 12:00 and 23:00), with the primary peak at 12:00 (7% of all suicides) and secondary peaks at 17:00 (6%) and 21:00 (6%). The lowest proportion of suicides occurred in the early morning at 05:00 (2%).

When individuals were stratified by BAL, different patterns emerged. The BAL<sub>H</sub> group had higher rates of suicide from 17:00 to 01:00, with a peak at 21:00. In contrast, the BAL<sub>O</sub> group had substantially higher rates of suicide than the BAL<sub>H</sub> group from 08:00 to 15:00, with a peak at around 12:00. The groups had similar rates of suicide between 02:00 and 07:00, with the nadir of suicide for BAL<sub>H</sub> and BAL<sub>O</sub> at 05:00 and 03:00, respectively.

Using a Bonferroni-adjusted significance level of .002, there were 6 time points at which the BAL<sub>H</sub> group ( $n = 2,222$ ) differed significantly from the BAL<sub>O</sub> group ( $n = 1,439$ ). When suicides occurred at 21:00 or 00:00, the individual was more likely to be in the BAL<sub>H</sub> category. In contrast, suicides at 08:00, 09:00, 12:00, or 14:00 were more likely to be in the BAL<sub>O</sub> category. Predictors of BAL<sub>H</sub> in individual bivariate analyses within each 4-hour time bin of the day are listed in Table 2.

### Predictors of BAL<sub>H</sub> Within Each 4-Hour Time Bin of the Day in Multivariable Analyses

**00:01–04:00.** Prior mental health treatment and suicide by poisoning were associated with decreased odds of having a BAL<sub>H</sub> at the time of death (Table 3).

**04:01–08:00.** Younger age and depressed mood around the time of suicide were associated with decreased odds of having a BAL<sub>H</sub> at the time of death.

**08:01–12:00.** Depressed mood around the time of suicide was associated with an increased probability of having a BAL<sub>H</sub> at the time of death.

**12:01–16:00.** A recent criminal legal problem was associated with decreased odds of having a BAL<sub>H</sub> at the time of death.

**16:01–20:00.** Being white was associated with nearly 2.5 times the odds of having a BAL<sub>H</sub> at the time of death compared to nonwhite individuals, whereas previous treatment for a mental health problem was associated with a decreased odds of having a BAL<sub>H</sub> at the time of death.

**20:01–0:00.** Individuals who were never married, veterans, and those with previous mental health treatment had a decreased probability of having a BAL<sub>H</sub> at the time of death. Having a problem at work and using a firearm to commit suicide were associated with an increased probability of having BAL<sub>H</sub> at the time of death.

### Weapons, Alcohol Consumption, and Circadian Pattern

The most common class of weapons used to commit suicide was handguns, followed by shotguns and rifles. However, no differences were observed for type, caliber, or other weapon-related variables when these were compared across alcohol consumption categories. When suicides were examined across circadian bins, gun ownership status was a significant predictor ( $P = .04$ ), with the victim being more likely to be the owner of the gun (Table 4).

### DISCUSSION

Despite evidence of a circadian pattern of suicides in the general population and both alcohol use and dependence being established risk factors for suicide, little is known about the circadian pattern of suicides in AD individuals. Here, we found that, overall, suicides peaked at 12:00, with a nadir at 05:00. A majority of individuals who committed suicide had a high BAL ( $\geq 80$  mg/dL) and a later peak time of completed suicide (21:00) than unintoxicated individuals (12:00). Intoxicated individuals who committed suicide later in the evening had more risk and protective factors than intoxicated individuals who committed suicide

It is illegal to post this copyrighted PDF on any website.

Table 2. Predictors of BAL<sub>H</sub> Within Each 4-Hour Time Bin of the Day in Individual Bivariate Analyses<sup>a</sup>

	00:01–04:00		04:01–08:00		08:01–12:00		12:01–16:00		16:01–20:00		20:01–00:00		p <sup>b</sup>
	(N = 495)	OR (95% CI)	(N = 375)	OR (95% CI)	(N = 660)	OR (95% CI)	(N = 657)	OR (95% CI)	(N = 755)	OR (95% CI)	(N = 719)	OR (95% CI)	
<b>Demographics</b>													
Age	37.3 (13.2)	0.99 (0.98–1.01)	41.5 (13.8)		45.4 (12.9)	0.99 (0.98–1.01)	44.8 (12.8)	1.01 (1.00–1.02)	44.4 (13.1)	1.02 (1.00–1.03)	41.4 (12.9)	1.00 (0.98–1.01)	<.0001
Male	86.1 (426)	0.91 (0.51–1.60)	83.7 (314)		84.2 (556)	0.97 (0.63–1.47)	80.4 (528)	1.22 (0.83–1.80)	84.9 (641)	1.39 (0.92–2.10)	86.4 (621)	1.13 (0.72–1.80)	.7277
White <sup>c</sup>	89.9 (445)	0.93 (0.49–1.79)	88.3 (331)		94.6 (624)	1.38 (0.69–2.75)	92.9 (610)	1.17 (0.65–2.13)	93.5 (706)	<b>2.41 (1.35–4.32)</b>	93.9 (675)	<b>1.91 (1.03–3.55)</b>	.0012
Hispanic	6.9 (34)	0.99 (0.46–2.12)	10.2 (38)		7.6 (50)	0.98 (0.50–1.93)	4.4 (29)	0.50 (0.23–1.09)	5.2 (39)	0.92 (0.46–1.82)	5.0 (36)	1.08 (0.51–2.29)	.0023
Marital status													.3097
Married	33.4 (165)	ref	37.8 (141)		38.5 (254)	ref	40.4 (264)	ref	42.9 (321)	ref	40.1 (287)	ref	
Never married	43.7 (216)	1.25 (0.80–1.96)	35.1 (131)		26.3 (173)	1.15 (0.78–1.70)	27.1 (177)	0.74 (0.51–1.09)	27.0 (202)	<b>0.59 (0.40–0.86)</b>	32.3 (231)	<b>0.52 (0.36–0.77)</b>	
Other <sup>d</sup>	22.9 (113)	1.07 (0.64–1.80)	27.1 (101)		35.2 (232)	1.13 (0.79–1.61)	32.5 (212)	0.99 (0.69–1.42)	30.1 (225)	0.89 (0.61–1.30)	27.7 (198)	0.84 (0.56–1.28)	
Veteran	13.5 (65)	0.91 (0.51–1.60)	19.4 (70)		19.6 (125)	1.08 (0.73–1.60)	19.7 (123)	1.01 (0.68–1.50)	21.0 (150)	1.19 (0.80–1.78)	22.6 (154)	<b>0.66 (0.45–0.97)</b>	.0004
Homeless	1.2 (6)	0.82 (0.15–4.52)	1.9 (7)		2.2 (14)	1.16 (0.40–3.34)	1.1 (7)	0.73 (0.16–3.29)	1.0 (7)	1.17 (0.23–6.06)	1.0 (7)	1.06 (0.20–5.51)	.1777
<b>Likely predisposing factors</b>													
Previous treatment for mental health problem	39.6 (196)	<b>0.46 (0.31–0.68)</b>	41.9 (157)		41.2 (318)	0.76 (0.56–1.03)	50.7 (333)	0.52 (0.39–0.72)	43.6 (329)	<b>0.67 (0.49–0.91)</b>	43.1 (310)	<b>0.54 (0.39–0.74)</b>	.3785
Current depressed mood	46.9 (232)	0.79 (0.54–1.16)	46.9 (176)		55.5 (366)	<b>0.62 (0.41–0.93)</b>	51.5 (338)	1.06 (0.78–1.43)	48.3 (365)	0.91 (0.67–1.23)	47.3 (340)	1.14 (0.82–1.57)	.6822
Previous suicide attempts	25.3 (125)	0.73 (0.47–1.12)	23.7 (89)		26.1 (172)	0.97 (0.60–1.56)	27.7 (182)	0.79 (0.56–1.11)	23.1 (174)	0.76 (0.53–1.09)	26.0 (187)	0.76 (0.53–1.09)	.9793
Physical health problem	12.1 (60)	0.74 (0.42–1.31)	14.9 (56)		15.3 (101)	<b>0.54 (0.31–0.97)</b>	16.9 (111)	0.88 (0.58–1.32)	16.2 (122)	1.07 (0.70–1.63)	14.2 (102)	0.73 (0.47–1.13)	.2804
Job problem	19.8 (98)	1.04 (0.64–1.70)	16.8 (63)		22.0 (145)	0.79 (0.46–1.35)	19.9 (131)	1.07 (0.73–1.57)	18.9 (143)	1.23 (0.82–1.84)	18.1 (130)	<b>1.83 (1.15–2.91)</b>	.5155
Recent criminal legal problem	13.5 (67)	1.26 (0.70–2.26)	14.1 (53)		21.2 (140)	0.73 (0.41–1.31)	16.4 (108)	<b>0.65 (0.43–0.99)</b>	14.0 (106)	0.76 (0.50–1.17)	14.7 (106)	1.26 (0.78–2.01)	.6282
Financial problem	17.8 (88)	1.13 (0.67–1.89)	14.4 (54)		17.3 (114)	0.59 (0.33–1.05)	16.4 (108)	0.85 (0.57–1.29)	15.8 (119)	0.89 (0.59–1.35)	14.6 (105)	1.24 (0.77–1.98)	.2308
<b>Suicide injury-related factors</b>													
Season of the year													.1120
Winter (Dec–Feb)	26.3 (130)	0.76 (0.43–1.32)	21.9 (82)		25.9 (171)	1.05 (0.58–1.91)	21.8 (143)	1.12 (0.72–1.72)	24.2 (183)	1.12 (0.72–1.72)	21.1 (152)	0.88 (0.55–1.41)	
Spring (March–May)	23.4 (116)	ref	25.6 (96)		20.3 (134)	ref	29.8 (196)	ref	24.4 (184)	ref	25.0 (180)	ref	
Summer (June–Aug)	27.7 (137)	0.76 (0.44–1.32)	25.1 (94)		31.1 (205)	0.75 (0.42–1.32)	26.6 (175)	1.17 (0.78–1.76)	27.4 (207)	1.17 (0.78–1.76)	28.9 (208)	1.35 (0.86–2.14)	
Fall (Sept–Nov)	22.6 (112)	0.91 (0.51–1.64)	27.5 (103)		22.7 (150)	1.00 (0.57–1.75)	21.8 (143)	1.44 (0.93–2.21)	24.0 (181)	1.44 (0.93–2.21)	24.9 (179)	0.75 (0.48–1.17)	
Weapon used													<.0001
Firearm	54.9 (270)	1.17 (0.56–2.48)	50.7 (190)		53.6 (353)	1.81 (0.84–3.87)	57.5 (378)	1.62 (0.92–2.85)	65.0 (491)	1.62 (0.92–2.85)	63.2 (454)	<b>2.50 (1.32–4.71)</b>	
Hanging, strangulation, suffocation	25.8 (127)	0.89 (0.40–1.96)	26.9 (101)		22.0 (145)	<b>2.32 (1.03–5.23)</b>	19.3 (127)	1.26 (0.67–2.37)	16.2 (122)	1.26 (0.67–2.37)	20.8 (149)	1.46 (0.74–2.90)	
Poisoning	11.2 (55)	<b>0.37 (0.15–0.88)</b>	13.9 (52)		18.4 (121)	1.84 (0.76–4.50)	14.6 (96)	0.54 (0.27–1.06)	11.4 (86)	0.54 (0.27–1.06)	9.9 (71)	0.78 (0.37–1.67)	
Other <sup>e</sup>	8.1 (40)	ref	8.5 (32)		6.1 (40)	ref	8.5 (56)	ref	7.4 (56)	ref	6.1 (44)	ref	

<sup>a</sup>Values expressed as % (n) unless otherwise indicated. Odds ratios computed with BAL<sub>0</sub> as the reference group. Bold indicates a significant bivariate association at  $\alpha = 0.05$ .

<sup>b</sup>p value corresponds to differences between time bin groups.

<sup>c</sup>Compared to a combined race category, which includes, but is not limited to, black, American Indian, Asian/Pacific Islander.

<sup>d</sup>Includes widowed, divorced, married but separated, and single (not otherwise specified).

<sup>e</sup>Includes, but is not limited to, non-powder gun, sharp instrument, blunt instrument, fall, explosive, drowning, fire or burns, motor vehicle (eg, buses, motorcycles), other transport vehicle (eg, trains, planes, boats).

Abbreviations: BAL<sub>H</sub> = heavy drinking (based on blood alcohol level), CI = confidence interval, OR = odds ratio, ref = reference group.

You are prohibited from making this PDF publicly available.

**Table 3. Predictors of BAL<sub>H</sub> Within Each 4-Hour Time Bin of the Day in Multivariable Analyses<sup>a</sup>**

Time	Variable	aOR (95% CI)
00:01–04:00	Previous treatment for mental health problem**	0.51 (0.34–0.77)
	Weapon used*	
	Firearm	1.05 (0.49–2.25)
	Poisoning	0.39 (0.16–0.93)
	Hanging, strangulation, suffocation	0.77 (0.34–1.71)
	Other <sup>b</sup>	ref
04:01–08:00	Age***	0.97 (0.95–0.98)
	Current depressed mood*	0.59 (0.39–0.90)
08:01–12:00	Current depressed mood*	1.46 (1.07–1.99)
12:01–16:00	Recent criminal legal problem*	0.65 (0.43–0.99)
16:01–20:00	White**	2.49 (1.38–4.47)
	Previous treatment for mental health problem**	0.66 (0.48–0.90)
20:01–00:00	Marital status**	
	Married	ref
	Never married	0.56 (0.37–0.83)
	Other <sup>c</sup>	0.94 (0.60–1.47)
	Veteran**	0.52 (0.35–0.79)
	Previous treatment for mental health problem*	0.63 (0.44–0.90)
	Job problem*	1.78 (1.08–2.93)
	Weapon used***	
	Firearm	2.47 (1.25–4.91)
	Poisoning	0.92 (0.41–2.06)
	Hanging, strangulation, suffocation	1.41 (0.68–2.95)
	Other <sup>b</sup>	ref

<sup>a</sup>P values correspond to Type III analysis of effects.<sup>b</sup>Includes, but is not limited to, non-powder gun, sharp instrument, blunt instrument, fall, explosive, drowning, fire or burns, motor vehicle (eg, buses, motorcycles), other transport vehicle (eg, trains, planes, boats).<sup>c</sup>Includes widowed, divorced, married but separated, and single (not otherwise specified).

\*P &lt; .05.

\*\*P &lt; .01.

\*\*\*P &lt; .001.

Abbreviations: aOR = adjusted odds ratio, BAL<sub>H</sub> = heavy drinking (based on blood alcohol level), ref = reference group.

at other times. Finally, intoxicated individuals who committed suicide in the afternoon or evening were more likely to do so with a gun.

We found that suicide was more likely to occur at midday in our AD sample, which is in line with prior community-based studies.<sup>25–27</sup> Suicides in intoxicated individuals peaked in the evening and were more likely to involve a gun, also consistent with prior findings<sup>33</sup> and with the notion that people are more likely to drink alcohol later in the day, which predisposes them to impulsive and suicidal behavior.<sup>33</sup> Our findings differ from those of Prasanna and colleagues, who found that suicide attempts in intoxicated AD individuals occurred most often in the morning (36.7%) and at night (26.7%).<sup>34</sup> The disparity between their study results and ours may be due to their small sample size (N = 30). Alternatively, lethal suicidal intent may be associated with different peak times than nonlethal intent.

Multiple factors influence the later time for the peak incidence of suicides in individuals who are intoxicated. Animal studies have shown a greater sensitivity to the depressant effects of alcohol around the usual sleep time,<sup>42,43</sup> which may lead to more unpredictable and impulsive behavior.<sup>33,44,45</sup> Heavy drinking may also be associated with mood disturbance, a sense of hopelessness,<sup>15,44,46</sup> and problematic interactions with family or friends, especially in the evening, when people usually socialize. Recurrent drinking, stress, and maladaptive behavior could lead

to a repetitive cycle from which suicide is viewed as an escape.<sup>15,47</sup>

Psychiatric disorders commonly co-occur with alcohol dependence,<sup>23</sup> and treating both disorders can decrease suicide risk in AD individuals.<sup>44</sup> The most common psychiatric disorders in individuals in this study were depressive disorder and dysthymic disorder. About half of the AD individuals had a depressed mood prior to suicide, a powerful risk factor for suicide in AD individuals with depressive disorder and heavy drinking.<sup>44</sup> Depressed mood may fluctuate across the day in distressed individuals, with the greatest intensity in the morning and a gradual improvement over the course of the day.<sup>48</sup> Thus, an association between depressed mood and BAL<sub>H</sub> in the 08:00–12:00 bin is not surprising.

However, the inverse relationship between depressed mood and BAL<sub>H</sub> in the 04:01–08:00 time bin may have been due to the interaction between sleep-related problems and a circadian variation in mood. That is, because interrupted sleep and early morning awakening are common symptoms of alcohol use and depressive disorder,<sup>49,50</sup> individuals could have awakened depressed, and, lacking access to alcohol, could not use it for self-medication.

Mental health treatment is another area in which some unique findings were obtained on post hoc analysis. Only 22% of the total sample and 13% of those with a history of psychiatric disorders screened positive for antidepressant medication on toxicologic testing. This may reflect nonadherence (with psychiatric care, psychotropic medications, or both) or a poor treatment response from the medication. In a recent meta-analytic review of data from 70 clinical trials of antidepressant medications,<sup>51</sup> investigators at the Nordic Cochrane Center found a lack of association between suicidal behavior and antidepressant medications in adults. It is possible that suicidal behavior emerging during treatment may be secondary to the worsening of mood and anxiety symptoms in the absence of a therapeutic response to an antidepressant medication rather than a direct adverse effect of it, as shown before.<sup>52</sup> Thus, improving access to psychiatric treatment and monitoring treatment compliance and effectiveness<sup>53</sup> could reduce the risk of suicide in this population.

The BAL<sub>O</sub> subgroup had a significantly higher proportion of individuals with prior mental health treatment (53% vs 40%), and individuals with positive toxicology screens for antidepressants were more likely to have a BAL<sub>O</sub> within the hours of 20:01 and 08:00. Thus, individuals receiving psychiatric treatment may have drunk less than those who needed treatment but were not receiving it. However, their conduct of suicide raises questions about other possible risk factors such as problematic social supports, comorbid medical disorders,<sup>21–23</sup> and possibly antidepressant medications themselves.<sup>54–56</sup>

**It is illegal to post this copyrighted PDF on any website.**

**Table 4. Weapon-Related Details and Circadian Bins<sup>a</sup>**

	00:01–04:00 (N=270)	04:01–08:00 (N=190)	08:01–12:00 (N=353)	12:01–16:00 (N=378)	16:01–20:00 (N=491)	20:01–00:00 (N=454)	P
Firearm type							.20
Handgun	176 (65.2)	128 (67.4)	235 (66.6)	267 (70.6)	342 (69.7)	324 (71.4)	
Shotgun	38 (14.1)	26 (13.7)	53 (15.0)	64 (16.9)	74 (15.1)	67 (14.8)	
Rifle	49 (18.1)	32 (16.8)	59 (16.7)	43 (11.4)	67 (13.6)	55 (12.1)	
Other	1 (0.4)	1 (0.5)	3 (0.8)	1 (0.3)	1 (0.2)	0 (0.0)	
Combination	0 (0.0)	1 (0.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Not reported	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Unknown	6 (2.2)	2 (1.1)	3 (0.8)	3 (0.8)	7 (1.4)	8 (1.8)	
Firearm caliber	N=187	N=143	N=266	N=273	N=345	N=336	.26
Mean (SD), in	.33 (.08)	.34 (.08)	.33 (.08)	.34 (.07)	.34 (.07)	.34 (.07)	
Median (IQR), in	.35 (.16)	.36 (.11)	.36 (.16)	.36 (.09)	.36 (.10)	.36 (.08)	
Range, in	.22–.50	.22–.54	.22–.50	.22–.46	.17–.45	.22–.50	
Stolen firearm							.54
No	36 (13.3)	35 (18.4)	48 (13.6)	44 (11.6)	58 (11.8)	55 (12.1)	
Yes	1 (0.4)	1 (0.5)	2 (0.6)	2 (0.5)	3 (0.6)	0 (0.0)	
Not applicable	7 (2.6)	2 (1.1)	6 (1.7)	10 (2.6)	10 (2.0)	14 (3.1)	
Unknown	226 (83.7)	152 (80.0)	297 (84.1)	322 (85.2)	420 (85.6)	385 (84.8)	
Stored gun locked							.66
Not locked	0 (0.0)	2 (1.1)	5 (1.4)	1 (0.3)	4 (0.8)	4 (0.9)	
Locked	2 (0.7)	0 (0.0)	2 (0.6)	3 (0.8)	3 (0.6)	3 (0.7)	
Other	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	
Not applicable	64 (23.7)	51 (26.8)	74 (21.0)	89 (23.5)	113 (23.0)	107 (23.6)	
Unknown	204 (75.5)	137 (72.1)	272 (77.1)	285 (75.4)	371 (75.5)	339 (74.6)	
Stored gun loaded							.28
Unloaded	0 (0.0)	0 (0.0)	3 (0.8)	2 (0.5)	0 (0.0)	4 (0.9)	
Loaded	0 (0.0)	1 (0.5)	2 (0.6)	0 (0.0)	1 (0.2)	2 (0.4)	
Not applicable	63 (23.3)	51 (26.8)	72 (20.4)	89 (23.5)	113 (23.0)	107 (23.6)	
Unknown	207 (76.6)	138 (72.6)	276 (78.2)	287 (75.9)	377 (76.7)	341 (75.1)	
Gun owner							.04
Shooter	30 (11.1)	15 (7.9)	39 (11.0)	29 (7.7)	45 (9.2)	36 (7.9)	
Parent	5 (1.9)	1 (0.5)	5 (1.4)	2 (0.5)	5 (1.0)	1 (0.2)	
Other family member	2 (0.7)	2 (1.1)	9 (2.5)	5 (1.3)	2 (0.4)	9 (2.0)	
Friend/acquaintance	4 (1.5)	6 (3.2)	6 (1.7)	6 (1.6)	4 (0.8)	11 (2.4)	
Stranger	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	0 (0.0)	
Other	0 (0.0)	0 (0.0)	1 (0.3)	1 (0.3)	7 (1.4)	3 (0.7)	
Unknown	228 (84.5)	166 (87.4)	293 (83.0)	335 (88.6)	427 (87.0)	394 (86.8)	

<sup>a</sup>Values expressed as n (%) unless otherwise indicated. Cochran-Mantel-Haenszel statistics used for categorical variables (row mean scores differ for the general association statistic for the circadian bin); analyses of variance used for the continuous variable (caliber).

Abbreviation: IQR = interquartile range.

Our finding that firearms were the most common means of suicide is highly consistent with both prior literature and recent CDC findings demonstrating this association.<sup>2</sup> The association between firearms and alcohol intoxication in the evening may reflect impulsivity augmented by intoxication, which in the absence of lethal means (eg, firearms) may not result in death. Other risk factors that were associated with alcohol intoxication at the time of death included being white and having job-related problems, established risk factors for both heavy drinking and suicidal behavior.<sup>4,5,57</sup> The association between current job problems and alcohol intoxication seen in the late evening hours may be a particularly relevant set of co-occurring risk factors, as seen previously.<sup>39</sup>

There are a number of limitations of the current investigation, including that the time of death was based on the time of injury, and these times are not always the same. Because the selection of an enriched sample of AD individuals with available BAL data may have introduced a selection bias, the sample may not be representative of all AD individuals who complete suicide. In addition, some

individuals may have been misclassified as to the presence of alcohol dependence. It is also possible that the BAL at the time of death does not reflect the individuals' habitual drinking patterns. The current employment status of most of the individuals was unavailable. Therefore, we used job-related problems at the time of death as an indicator that an individual was employed and as a measure of work-associated problems. However, we do not know how correlated current employment status and job-related problems were. Intoxicated individuals had relatively fewer health problems and past suicide attempts but more job-related problems. Although comorbid mood and anxiety disorders are prevalent in alcohol dependence,<sup>23</sup> fewer medical and psychiatric problems identified in intoxicated individuals may reflect an ascertainment bias, or the lack of an adequate clinical history. The toxicologic evidence of antidepressant medications may have been an underestimate due to partial or complete nonadherence with these medications,<sup>58,59</sup> and the duration of antidepressant medication use is unclear for individuals who tested positive for them. Finally, we were unable to adjust for the proportion of AD individuals awake



at each hour<sup>60</sup> and unable to assess how different factors such as social isolation, occupational problems, alcohol consumption, and access to weapons acted either alone or in combination to increase the risk of suicide.

Several directions are available for future research. Considering the novel findings in this study, an attempt should be made to replicate these findings. Future studies should also account for the role of associated factors such as sleep continuity disturbance, circadian patterns of alcohol metabolism, and executive functioning and how

psychological stress and social problems interact with alcohol use to increase risk of suicide.

## CONCLUSION

The peak incidence of suicide among AD individuals was later in the day for those who were intoxicated than those with moderate or no recent alcohol consumption. In addition, intoxication in the evening may represent a greater risk for suicide, particularly using lethal means such as firearms.

**Submitted:** July 9, 2017; accepted April 24, 2018.

**Published online:** October 23, 2018.

**Potential conflicts of interest:** Dr Chakravorty has received research support from AstraZeneca and Teva. Dr Perlis has received research support from Nexalin Technology and Teva. Dr Grandner has received research support from Nexalin technology and Kemin Industries and serves as a consultant for Fitbit, Curaegis Technologies, and Natrol. Dr Kranzler has served as a consultant, advisory board member, or continuing medical education speaker for Alkermes, Indivior, and Lundbeck and is also a member of the American Society of Clinical Psychopharmacology Clinical Trials Initiative (ACTIVE) group, which was supported in the last 3 years by AbbVie, Alkermes, Amygdala Neurosciences, Arbor, Ethypharm, Indivior, Lilly, Lundbeck, Otsuka, and Pfizer. Dr Smith certifies that she has no actual or potential conflict of interest with this investigation.

**Funding/support:** The study was not supported by any independent grant funding. Partial salary support was provided by the following grants: Department of Veterans Affairs grant IK2CX000855 (Dr Chakravorty), National Institutes of Health (NIH) grants K23 HL110216 and NIH R21 ES022931 (Dr Grandner); R01 AG041783 (Dr Perlis) and R56 AG050620 (Dr Perlis); R01 AA023192 (Dr Kranzler) and R01 AA021164 (Dr Kranzler).

**Role of sponsor:** None.

**Disclaimer:** The content of this publication does not represent the views of the University of Pennsylvania, Department of Veterans Affairs, the United States Government, or any other institution.

**Previous presentation:** Some of the results from this study were presented at the SLEEP 2016 annual meeting; June 11–15, 2016; in Denver, Colorado.

**Acknowledgments:** The authors thank the Centers for Disease Control and Prevention, Atlanta, Georgia, for providing access to the NVDRS dataset. They also express gratitude to the following individuals who helped with the data analysis and provided intellectual assistance with the draft: Ninad Chaudhary, MBBS, MPH, University of Alabama at Birmingham; and Kachina Allen, PhD, Princeton University. Neither Dr Chaudhary nor Dr Allen has any conflict of interest to report with the manuscript draft.

## REFERENCES

- Centers for Disease Control and Prevention (CDC). Preventing suicide. CDC website. <http://www.cdc.gov/Features/PreventingSuicide/index.html>. Accessed May 15, 2017.
- Curtin SC, Warner M, Hedegaard H. Increase in suicide in the United States, 1999–2014. *NCHS Data Brief*. 2016;(241):1–8.
- Crump C, Sundquist K, Sundquist J, et al. Sociodemographic, psychiatric and somatic risk factors for suicide: a Swedish national cohort study. *Psychol Med*. 2014;44(2):279–289.
- Heikkinen ME, Isometsä ET, Marttunen MJ, et al. Social factors in suicide. *Br J Psychiatry*. 1995;167(6):747–753.
- Chang SS, Stuckler D, Yip P, et al. Impact of 2008 global economic crisis on suicide: time trend study in 54 countries. *BMJ*. 2013;347:f5239.
- Milner A, Spittal MJ, Pirkis J, et al. Suicide by occupation: systematic review and meta-analysis. *Br J Psychiatry*. 2013;203(6):409–416.
- Haukka J, Suominen K, Partonen T, et al. Determinants and outcomes of serious attempted suicide: a nationwide study in Finland, 1996–2003. *Am J Epidemiol*. 2008;167(10):1155–1163.
- Pokorny AD. Prediction of suicide in psychiatric patients: report of a prospective study. *Arch Gen Psychiatry*. 1983;40(3):249–257.
- Kontaxakis VP, Christodoulou GN, Mavreas VG, et al. Attempted suicide in psychiatric outpatients with concurrent physical illness. *Psychother Psychosom*. 1988;50(4):201–206.
- Juurink DN, Herrmann N, Szalai JP, et al. Medical illness and the risk of suicide in the elderly. *Arch Intern Med*. 2004;164(11):1179–1184.
- Helmkamp JC. Occupation and suicide among males in the US Armed Forces. *Ann Epidemiol*. 1996;6(1):83–88.
- Tidemalm D, Långström N, Lichtenstein P, et al. Risk of suicide after suicide attempt according to coexisting psychiatric disorder: Swedish cohort study with long term follow-up. *BMJ*. 2008;337:a2205.
- Mukamal KJ, Kawachi I, Miller M, et al. Drinking frequency and quantity and risk of suicide among men. *Soc Psychiatry Psychiatr Epidemiol*. 2007;42(2):153–160.
- Centers for Disease Control and Prevention (CDC). Toxicology testing and results for suicide victims—13 states, 2004. *MMWR Morb Mortal Wkly Rep*. 2006;55(46):1245–1248.
- Berglund M, Ojehagen A. The influence of alcohol drinking and alcohol use disorders on psychiatric disorders and suicidal behavior. *Alcohol Clin Exp Res*. 1998;22(suppl):333S–345S.
- Klatsky AL, Armstrong MA. Alcohol use, other traits, and risk of unnatural death: a prospective study. *Alcohol Clin Exp Res*. 1993;17(6):1156–1162.
- Andreasson S, Allebeck P, Romelsjö A. Alcohol and mortality among young men: longitudinal study of Swedish conscripts. *Br Med J (Clin Res Ed)*. 1988;296(6628):1021–1025.
- Borges G, Cherpitel CJ, MacDonald S, et al. A case-crossover study of acute alcohol use and suicide attempt. *J Stud Alcohol*. 2004;65(6):708–714.
- Hedden SL, Kennet J, Lipari R, et al. *Behavioral Health Trends in the United States: Results from the 2014 National Survey on Drug Use and Health*. Rockville, MD: SAMHSA; 2015.
- Murphy GE, Wetzell RD. The lifetime risk of suicide in alcoholism. *Arch Gen Psychiatry*. 1990;47(4):383–392.
- Miller WR, Tonigan JS, Longabaugh R. The Drinker Inventory of Consequences (DrlNC): An Instrument for Assessing Adverse Consequences of Alcohol Abuse. National Institute on Alcohol Abuse and Alcoholism Project MATCH Monograph Series, 1995. National Institutes of Health website. <http://pubs.niaaa.nih.gov/publications/ProjectMatch/match04.pdf>. Accessed May 15, 2017.
- Mannelli P, Pae CU. Medical comorbidity and alcohol dependence. *Curr Psychiatry Rep*. 2007;9(3):217–224.
- Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2004;61(8):807–816.
- Williams P, Tansella M. The time for suicide. *Acta Psychiatr Scand*. 1987;75(5):532–535.
- Altamura C, VanGastel A, Pioli R, et al. Seasonal and circadian rhythms in suicide in Cagliari, Italy. *J Affect Disord*. 1999;53(1):77–85.
- Preti A, Miotto P. Diurnal variations in suicide by age and gender in Italy. *J Affect Disord*. 2001;65(3):253–261.
- van Houwelingen CA, Beersma DG. Seasonal changes in 24-h patterns of suicide rates: a study on train suicides in The Netherlands. *J Affect Disord*. 2001;66(2–3):215–223.
- Erazo N, Baumert J, Ladwig KH. Sex-specific time patterns of suicidal acts on the German railway system: an analysis of 4003 cases. *J Affect Disord*. 2004;83(1):1–9.
- Vollen KH, Watson CG. Suicide in relation to time of day, day of week. *Am J Nurs*. 1975;75(2):263.
- Gallerani M, Avato FM, Dal Monte D, et al. The time for suicide. *Psychol Med*. 1996;26(4):867–870.
- Barracough BM. Time of day chosen for suicide. *Psychol Med*. 1976;6(2):303–305.
- Maldonado G, Kraus JF. Variation in suicide occurrence by time of day, day of the week, month, and lunar phase. *Suicide Life Threat Behav*. 1991;21(2):174–187.
- Welte JW, Abel EL, Wiczorek W. The role of alcohol in suicides in Erie County, NY, 1972–84. *Public Health Rep*. 1988;103(6):648–652.
- Prasanna CA, Gandhibabu R, Asok Kumar M. Risk factors in suicide among male alcohol dependents. *Journal of Evolution of Medical and Dental Sciences*. 2014;3(52):12073–12088.
- Center for Disease Control and Prevention (CDC). National Violent Death Reporting system (NVDRS). Violence Prevention, 2013. <https://www.cdc.gov/violenceprevention/nvdrs/>. Accessed May 15, 2017.
- Hempstead KA, Phillips JA. Rising suicide among adults aged 40–64 years: the role of job and financial circumstances. *Am J Prev Med*. 2015;48(5):491–500.



It is illegal to post this copyrighted PDF on any website.

37. National Institute on Alcohol Abuse and Alcoholism. Drinking Levels Defined. National Institutes of Health website. <http://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking>, 2017.
38. Lange JE, Voas RB. Defining binge drinking quantities through resulting BACs. *Annu Proc Assoc Adv Automot Med*. 2000;44:389–404.
39. Caetano R, Kaplan MS, Huguet N, et al. Precipitating circumstances of suicide and alcohol intoxication among US ethnic groups. *Alcohol Clin Exp Res*. 2015;39(8):1510–1517.
40. NVDRS. *National Violent Death Reporting System (NVDRS): Guidelines for data users*. Atlanta, GA: CDC; 2013:1–225.
41. SAS [computer program]. Version 9.4 for Windows. Cary, NC: SAS Institute Inc.
42. Brick J, Pohorecky LA, Faulkner W, et al. Circadian variations in behavioral and biological sensitivity to ethanol. *Alcohol Clin Exp Res*. 1984;8(2):204–211.
43. Sturtevant RP, Garber SL. Circadian rhythm of blood ethanol clearance rates in rats: response to reversal of the L/D regimen and to continuous darkness and continuous illumination. *Chronobiol Int*. 1988;5(2):137–148.
44. Sher L. Alcohol consumption and suicide. *QJM*. 2006;99(1):57–61.
45. Danel T, Touitou Y. Chronobiology of alcohol: from chronokinetics to alcohol-related alterations of the circadian system. *Chronobiol Int*. 2004;21(6):923–935.
46. Sher L. Risk and protective factors for suicide in patients with alcoholism. *Sci World J*. 2006;6:1405–1411.
47. Preuss UW, Schuckit MA, Smith TL, et al. Comparison of 3190 alcohol-dependent individuals with and without suicide attempts. *Alcohol Clin Exp Res*. 2002;26(4):471–477.
48. Daly M, Delaney L, Doran PP, et al. The role of awakening cortisol and psychological distress in diurnal variations in affect: a day reconstruction study. *Emotion*. 2011;11(3):524–532.
49. Arnedt JT, Rohsenow DJ, Almeida AB, et al. Sleep following alcohol intoxication in healthy, young adults: effects of sex and family history of alcoholism. *Alcohol Clin Exp Res*. 2011;35(5):870–878.
50. Murphy MJ, Peterson MJ. Sleep Disturbances in Depression. *Sleep Med Clin*. 2015;10(1):17–23.
51. Sharma T, Guski LS, Freund N, et al. Suicidality and aggression during antidepressant treatment: systematic review and meta-analyses based on clinical study reports. *BMJ*. 2016;352:i65.
52. Courtet P, Jaussent I, Lopez-Castroman J, et al. Poor response to antidepressants predicts new suicidal ideas and behavior in depressed outpatients. *Eur Neuropsychopharmacol*. 2014;24(10):1650–1658.
53. Zalsman G, Hawton K, Wasserman D, et al. Suicide prevention strategies revisited: 10-year systematic review. *Lancet Psychiatry*. 2016;3(7):646–659.
54. Brent DA. Antidepressants and suicidality. *Psychiatr Clin North Am*. 2016;39(3):503–512.
55. Braun C, Bschor T, Franklin J, et al. Suicides and suicide attempts during long-term treatment with antidepressants: a meta-analysis of 29 placebo-controlled studies including 6,934 patients with major depressive disorder. *Psychother Psychosom*. 2016;85(3):171–179.
56. Bielefeldt AO, Danborg PB, Göttsche PC. Precursors to suicidality and violence on antidepressants: systematic review of trials in adult healthy volunteers. *J R Soc Med*. 2016;109(10):381–392.
57. Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proc Natl Acad Sci U S A*. 2015;112(49):15078–15083.
58. Isacsson G, Bergman U, Rich CL. Antidepressants, depression and suicide: an analysis of the San Diego study. *J Affect Disord*. 1994;32(4):277–286.
59. Marzuk PM, Tardiff K, Leon AC, et al. Use of prescription psychotropic drugs among suicide victims in New York City. *Am J Psychiatry*. 1995;152(10):1520–1522.
60. Perlis ML, Grandner MA, Brown GK, et al. Nocturnal wakefulness as a previously unrecognized risk factor for suicide. *J Clin Psychiatry*. 2016;77(6):e726–e733.

**Editor's Note:** We encourage authors to submit papers for consideration as a part of our Focus on Suicide section. Please contact Philippe Courtet, MD, PhD, at [pcourtet@psychiatrist.com](mailto:pcourtet@psychiatrist.com).

You are prohibited from making this PDF publicly available.