

It is illegal to post this copyrighted PDF on any website.

# Paradoxical Trend for Improvement in Mental Health With Aging: A Community-Based Study of 1,546 Adults Aged 21–100 Years

Michael L. Thomas, PhD<sup>a</sup>; Christopher N. Kaufmann, PhD, MHS<sup>a,c</sup>; Barton W. Palmer, PhD<sup>a,c,d</sup>; Colin A. Depp, PhD<sup>a,c,d</sup>; Averria Sirkin Martin, PhD<sup>a,c</sup>; Danielle K. Gloriosi, MSW<sup>a,c</sup>; Wesley K. Thompson, PhD<sup>a,c</sup>; and Dilip V. Jeste, MD<sup>a,b,c,\*</sup>

## ABSTRACT

**Objective:** Studies of aging usually focus on trajectories of physical and cognitive function, with far less emphasis on overall mental health, despite its impact on general health and mortality. This study examined linear and nonlinear trends of physical, cognitive, and mental health over the entire adult lifespan.

**Methods:** Cross-sectional data were obtained from 1,546 individuals aged 21–100 years, selected using random digit dialing for the Successful Aging Evaluation (SAGE) study, a structured multicohort investigation that included telephone interviews and in-home surveys of community-based adults without dementia. Data were collected from 1/26/2010 to 10/07/2011 targeting participants aged 50–100 years and from 6/25/2012 to 7/15/2013 targeting participants aged 21–100 years with an emphasis on adding younger individuals. Data included self-report measures of physical health, measures of both positive and negative attributes of mental health, and a phone interview–based measure of cognition.

**Results:** Comparison of age cohorts using polynomial regression suggested a possible accelerated deterioration in physical and cognitive functioning, averaging 1.5 to 2 standard deviations over the adult lifespan. In contrast, there appeared to be a linear improvement of about 1 standard deviation in various attributes of mental health over the same life period.

**Conclusions:** These cross-sectional findings suggest the possibility of a linear improvement in mental health beginning in young adulthood rather than a U-shaped curve reported in some prior studies. Lifespan research combining psychosocial and biological markers may improve our understanding of resilience to mental disability in older age and lead to broad-based interventions promoting mental health in all age groups.

*J Clin Psychiatry* 2016;77(8):e1019–e1025

[dx.doi.org/10.4088/JCP.16m10671](https://doi.org/10.4088/JCP.16m10671)

© Copyright 2016 Physicians Postgraduate Press, Inc.

<sup>a</sup>Departments of Psychiatry and <sup>b</sup>Neurosciences and <sup>c</sup>Sam and Rose Stein Institute for Research on Aging, University of California, San Diego, La Jolla

<sup>d</sup>Veterans Affairs San Diego Healthcare System, San Diego, California

\*Corresponding author: Dilip V. Jeste, MD, Stein Institute for Research on Aging, University of California, San Diego, 9500 Gilman Dr, #0664, La Jolla, CA 92093-0738 ([djeste@ucsd.edu](mailto:djeste@ucsd.edu)).

Traditionally, aging has been conceptualized as an ongoing process of physical and cognitive decline, with far less emphasis on mental health. Yet, in a recent national poll, 89% of American adults reported that mental health was as important as physical health.<sup>1</sup> Impaired mental health can adversely impact self-care and is a leading cause of disability.<sup>2</sup> Furthermore, untreated mental health problems often have a deleterious effect on physical health.<sup>3</sup>

Although mental health of older people might be assumed to follow a deteriorating course in step with worsening physical and cognitive function, findings regarding subjective well-being have been mixed.<sup>4,5</sup> Some investigators find a U-shaped curve of well-being across the lifespan, with declines from early adulthood to middle age followed by improvement during later adulthood.<sup>6–8</sup> Other studies report contradictory findings,<sup>9–11</sup> including increase, decline, or flat trajectory of psychological well-being across early adulthood. Charles et al<sup>10</sup> showed declines in negative affect while positive affect remained stable. Blazer<sup>12</sup> and Vaillant,<sup>13</sup> among others, have described successful cognitive and emotional aging in specific groups of older adults.<sup>14–16</sup> The reasons for differences in results are not obvious. Springer et al<sup>17</sup> suggest that changes in well-being with age may be due to measurement variation across studies, with many studies focusing on single domains of physical<sup>18,19</sup> or cognitive health<sup>20</sup> or on single indicators of mental health such as psychological well-being.

Psychological well-being<sup>21</sup> should be distinguished from the multidimensional construct of mental health, which also includes perceived stress and absence of anxiety and depressive symptoms. The effects of age may vary depending on the specific domain(s) assessed. In one study,<sup>7</sup> happiness, enjoyment, and the inverse of sadness were all U-shaped, while stress, worry, and anger showed apparent improvements with aging.<sup>7</sup> Thus, the empirical literature is inconsistent regarding the pattern of change in mental health across the adult lifespan.

Epidemiologic studies have reported lower prevalence of all mental illnesses, except for neurocognitive disorders, in older adults.<sup>22–25</sup> The reasons remain unclear. Common explanations include inappropriate diagnostic criteria for geriatric mental disorders, older patients' reluctance or inability (due to memory difficulties) to report mental problems, and misattribution of psychological symptoms to physical illnesses.<sup>26</sup> However, survivor effects also may play a role—ie, the adults who survive into older age may have better mental health than younger counterparts. Additional studies of aging, using validated measures of broadly defined mental health along with physical and cognitive functioning across the adult lifespan, are needed.

We simultaneously examined physical health, cognitive function, and several measures of mental health across the adult lifespan (age

- Studies show declining physical and cognitive health during later phases of the lifespan; yet, few studies have examined trajectories of mental health in the context of these declines.
- Our results point to a progressive improvement in mental health (ie, higher levels of life satisfaction, happiness, and well-being and lower levels of anxiety, depression, and perceived stress) from age 21 years through the 90s.
- When treating younger adults, consider how to leverage positive mental health factors in augmenting their psychosocial functioning, whereas in older adults, consider how these factors may help with physical and cognitive functioning.

21–100 years) in a community sample. Unlike many studies using samples of convenience, we obtained a sample in San Diego County, California, using random digit dialing, with nearly equal numbers of men and women, stratified by age decade, and an oversampling of adults over age 75. We hypothesized that, while physical and cognitive function would be lower among successively older age cohorts, mental health would follow a nonlinear, U-shaped trend, with middle-aged cohorts evidencing the worst mental health relative to younger and older cohorts.

## METHODS

### Participants

Participants included 1,546 community-dwelling adults from the University of California, San Diego, Successful AGing Evaluation (SAGE) study.<sup>27</sup> A structured multicohort design was used to recruit residents of San Diego County. The study originally targeted adults aged 50–100 years (data collected from 1/26/2010 to 10/07/2011), but we subsequently targeted participants aged 21–100 years (data collected from 6/25/2012 to 7/15/2013) with an emphasis on adding younger individuals. Partial data in adults aged 50+ years have been reported previously<sup>27,28</sup> in regard to self-perceived successful aging, but this current report represents our first examination of the life-span trajectory of physical, cognitive, and mental health in the entire age range. Exclusion criteria were (1) residence in a nursing home or need for daily skilled nursing care, (2) self-reported prior diagnosis of dementia, (3) terminal illness or need for hospice care, and (4) inability to understand written or spoken English. Participants who completed a 25-minute structured telephone interview that included cognitive assessment (see Supplementary Text), were subsequently mailed a survey questionnaire that asked for detailed demographic and clinical information and included a number of rating scales and other measures. The enrollment sequence is depicted in Supplementary eFigure 1. Participants were compensated \$10 for their participation in the phone interview and \$15 for the completion of their baseline SAGE survey. The study was approved by the University of California, San Diego, Human Research Protections Program.

### Measures

**Physical health.** The construct of physical health was operationally defined as a composite of self-reported physical functioning, role-physical, bodily pain, and general health subscale total scores from the Medical Outcomes Study 36-item Short-Form Health Survey (SF-36).<sup>29,30</sup> Subjects rated themselves on both the presence of good (or poor) health and its functional consequences. To create the composite, we standardized scores from each subscale, computed the average value, and then standardized the outcome (z-scores). Unlike the manual-based scoring method for the SF-36, which includes both physical and mental health subscale scores to determine the physical health composite, our composite was composed of physical health subscales only, in order to operationalize physical health independently of mental health. The internal consistency of the modified physical composite was high ( $\alpha = .81$ ), and variance was dominated by a general factor ( $\omega_h = 0.75$ ).

**Cognitive function.** The construct of cognitive function was assessed by total scores on the modified Telephone Interview of Cognitive Status (TICS-M).<sup>31</sup> TICS-M scores are documented as having good convergent<sup>31</sup> and discriminant<sup>32</sup> validity.

**Mental health.** The construct of mental health was operationally defined as a composite of 3 measures reflecting positive and 3 measures reflecting negative attributes. Measures reflecting positive attributes included total scores on the happiness subscale<sup>33</sup> of the Center for Epidemiologic Studies-Depression scale (CES-D); the Satisfaction With Life Scale<sup>34</sup>; and a composite of vitality, social functioning, role-emotional, and mental health subscale total scores from the SF-36.<sup>29</sup> Negative attributes were assessed with total scores on the Brief Symptom Inventory (BSI) Anxiety scale,<sup>35</sup> Patient Health Questionnaire depression module (PHQ-9),<sup>36</sup> and Perceived Stress Scale.<sup>37,38</sup> We created the mental health composite by standardizing the scores reflecting 3 positive and 3 negative psychological attributes, calculating the average value (with negative attributes weighted by  $-1$ ) and then standardizing the outcome. The internal consistency of the composite was high ( $\alpha = .87$ ), and variance was dominated by a general factor ( $\omega_h = 0.82$ ).

### Analyses

To evaluate aging-associated trends in physical, cognitive, and mental health, we fitted power polynomial regression models to each construct.<sup>39</sup> These models sought to determine whether, and how, the relationship between age and each construct was linear or nonlinear. We hierarchically compared models that included linear, linear and quadratic, or linear, quadratic, and cubic effects of (centered) age. Polynomial terms allow nonlinear curves or bends in the relationship between variables. Models with linear terms allow zero bends, models with linear and a single nonlinear (quadratic) term allow 1 bend, and models with linear and 2 nonlinear (quadratic and cubic) terms allow 2 bends. The linear term indicates whether the relationship between

Table 1. Results of Nested Regression Models for Physical Health (N = 1,491<sup>a</sup>)

Predictor	Model 1		Model 2		Model 3		Model 4	
	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE
Intercept	0.0037	0.0252	−0.0004	0.0236	0.0638	0.0365	0.0769*	0.0380
Gender (male)	−0.0234	0.0534	0.0690	0.0503	0.0767	0.0504	0.0747	0.0504
Education	0.2398**	0.0406	0.1745**	0.0382	0.1682**	0.0383	0.1713**	0.0383
Marital status (married)	0.2816**	0.0539	0.2201**	0.0506	0.1831**	0.0531	0.1877**	0.0532
Age (linear)	...	...	−0.0166**	0.0011	−0.0182**	0.0013	−0.0155**	0.0025
Age (quadratic)	...	...	...	...	−0.0001*	0.0001	−0.0002*	0.0001
Age (cubic)	...	...	...	...	...	...	<.0001	<.0001
LRT <i>P</i> value	...	...	<.001	...	.021	...	.219	...
<i>R</i> <sup>2</sup>	0.051	...	0.170	...	0.173	...	0.174	...

<sup>a</sup>N = 1,491 reflects the sample size after deletion for missing data.\**P* < .05.\*\**P* < .001.

Symbol: ... = not applicable.

Abbreviations: *b* = regression parameter estimate, LRT = likelihood ratio test, *R*<sup>2</sup> = variance accounted for, SE = standard error.Table 2. Results of Nested Regression Models for Cognitive Function (N = 1,491<sup>a</sup>)

Predictor	Model 1		Model 2		Model 3		Model 4	
	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE
Intercept	0.0118	0.0244	0.0069	0.0219	0.1893***	0.0333	0.2114***	0.0347
Gender (male)	−0.3633***	0.0517	−0.2509***	0.0467	−0.2290***	0.0460	−0.2324***	0.0460
Education	0.3933***	0.0393	0.3138***	0.0355	0.2960***	0.0350	0.3011***	0.0350
Marital status (married)	0.2245***	0.0522	0.1498**	0.0470	0.0445	0.0485	0.0522	0.0485
Age (linear)	...	...	−0.0202***	0.0011	−0.0246***	0.0012	−0.0202***	0.0023
Age (quadratic)	...	...	...	...	−0.0004***	0.0001	−0.0005***	0.0001
Age (cubic)	...	...	...	...	...	...	<.0001*	<.0001
LRT <i>P</i> value	...	...	<.001	...	<.001	...	.024	...
<i>R</i> <sup>2</sup>	0.094	...	0.273	...	0.297	...	0.299	...

<sup>a</sup>N = 1,491 reflects the sample size after deletion for missing data.\**P* < .05.\*\**P* < .01.\*\*\**P* < .001.

Symbol: ... = not applicable.

Abbreviations: *b* = regression parameter estimate, LRT = likelihood ratio test, *R*<sup>2</sup> = variance accounted for, SE = standard error.

variables generally rises or falls; the nonlinear (quadratic and cubic) terms indicate whether the bends are concave upward or downward. All models also included gender and marital status (contrast coded) and education (linear trend coded).<sup>40</sup> Additionally, we assessed age effects for each scale contributing to the mental health composite. Models were compared using likelihood ratio tests. Effect size was quantified in terms of increment in variance accounted for ( $\Delta R^2$ ) by comparing nested models. All analyses were conducted using R.<sup>41</sup>

## RESULTS

### Participant Characteristics

Mean age was 66 years (standard deviation [SD] = 21 years), and 51% of the participants were men. Highest level of education completed was ≤ 12 years for 20%, 13–16 years (bachelor's degree) for 60%, and postgraduate education for 21% of the sample. Seventy-six percent identified as non-Latino white, 14% as Hispanic/Latino, 7% Asian American, 1% African American, and 2% other ethnic or racial backgrounds (see Supplementary eTable 1).

### Age-Associated Trends in Physical Health, Cognitive Function, and Mental Health

Tables 1 through 3 show results of regression analyses for physical, cognitive, and mental health variables, respectively. For ease of communication, results are described in terms of age-associated deterioration or improvement in health; however, within-subject changes can be inferred only from longitudinal data. Interpretive limitations of the cross-sectional design are discussed below.

While physical and cognitive function showed nonlinear (accelerating) worsening with age, mental health showed linear improvement. Specifically, the physical health model with gender, education, and marital status was improved by adding a linear effect of age ( $\Delta R^2 = 0.125$ ) as well as a nonlinear (quadratic) effect ( $\Delta R^2 = 0.004$ ), but not a second nonlinear (cubic) effect. The cognitive function model was improved by adding a linear effect of age ( $\Delta R^2 = 0.197$ ) as well as a nonlinear (quadratic) effect ( $\Delta R^2 = 0.033$ ) and a second nonlinear (cubic) effect ( $\Delta R^2 = 0.003$ ). The mental health model was improved by adding a linear effect of age ( $\Delta R^2 = 0.046$ ), but not by adding nonlinear (quadratic and cubic) effects (see Supplementary Text).

Table 3. Results of Nested Regression Models for Mental Health (N = 1,491<sup>a</sup>)

Predictor	Model 1		Model 2		Model 3		Model 4	
	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE
Intercept	−0.0003	0.0255	0.0022	0.0249	0.0483	0.0386	0.0651	0.0402
Gender (male)	0.0920	0.0540	0.0354	0.0532	0.0409	0.0533	0.0384	0.0533
Education	0.1859**	0.0411	0.2258**	0.0404	0.2213**	0.0405	0.2252**	0.0406
Marital status (married)	0.2038**	0.0546	0.2414**	0.0535	0.2149**	0.0562	0.2208**	0.0563
Age (linear)	...	...	0.0102**	0.0012	0.0090**	0.0014	0.0124**	0.0027
Age (quadratic)	...	...	...	...	−0.0001	0.0001	−0.0002*	0.0001
Age (cubic)	...	...	...	...	...	...	<.0001	<.0001
LRT <i>P</i> value	...	...	<.001	...	.119	...	.138	...
<i>R</i> <sup>2</sup>	0.036		0.080		0.082		0.083	

<sup>a</sup>N = 1,491 reflects the sample size after deletion for missing data.\**P* < .05.\*\**P* < .001.

Symbol: ... = not applicable.

Abbreviations: *b* = regression parameter estimate, LRT = likelihood ratio test, *R*<sup>2</sup> = variance accounted for, SE = standard error.

Figure 1. Predicted Values of Each Health Domain by Age





Figure 2. Predicted Values for Individual Measures of Mental Health by Age



Abbreviations: BSI = Brief Symptom Inventory, CES-D = Center for Epidemiologic Studies-Depression scale, PHQ-9 = Patient Health Questionnaire, SF-36 = Medical Outcomes Study 36-item Short-Form Health Survey.

Predicted values for the best fitting regression models are plotted in Figure 1 (corresponding observed data are reported in Supplementary eFigure 2). Successively older cohorts had better mental health despite having successively worse physical and cognitive function than younger cohorts. Across the 80 cross-sectional years of data, participants reported, on average, approximately a 1.5 SD drop in self-reported physical health and a 2 SD decline in cognitive function. Cognitive function appeared very nonlinear, with accelerating decline beginning in the sixties. The apparent improvement in mental health with age was linear, with participants reporting nearly 1 SD of change, on average.

It has been argued that the U-shaped curve frequently seen in studies of well-being is an artifact of inappropriately controlling for marital status and education, as these variables are possibly influenced by well-being.<sup>42</sup> In the present data, results without these covariates were similar except for a few minor differences—ie, the nonlinear (quadratic) effect of age on mental health was additionally significant and the second nonlinear (cubic) effect of age on cognitive function was nonsignificant. Interpretively, mental health showed a slight inverted U-shaped trend with age that peaked in late life, and cognitive function showed an inverted U-shaped trend that peaked in early life.

Figure 2 shows predicted age-related trends in the individual mental health scores that contributed to the composite. Scores on the happiness subscale (from the CES-D), and reverse-coded Anxiety scale (from the BSI) and depression module (PHQ-9) all improved with age (linear terms of *b* = 0.0108, standard error [SE] = 0.0013, *P* < .001; *b* = 0.0068, SE = 0.0013, *P* < .001; and *b* = 0.0085, SE = 0.0013, *P* < .001, respectively). Both the SF-36 mental health composite and the reverse-coded Perceived Stress Scale score demonstrated inverted U-shaped effects (linear terms of *b* = 0.0011, SE = 0.0015, *P* = .486 and *b* = 0.0075, SE = 0.0015, *P* < .001, respectively, and quadratic



**It is illegal to post this copyrighted PDF on any website.**

terms of  $b = -0.0002$ ,  $SE = 0.0001$ ,  $P = .005$  and  $b = -0.0001$ ,  $SE = 0.0001$ ,  $P = .037$ , respectively). The Satisfaction With Life Scale score demonstrated a U-shaped effect (linear term of  $b = 0.0155$ ,  $SE = 0.0014$ ,  $P < .001$  and quadratic term of  $b = 0.0001$ ,  $SE = 0.0001$ ,  $P < .029$ ).

## DISCUSSION

Our findings support the existence of a “paradox” in which aging is associated with better mental health among older adults at the population level despite loss of physical and cognitive function. Consistent with prior research, relative to successively younger cohorts, older cohorts had worse physical and cognitive function,<sup>43–45</sup> but better mental health. This pattern was observed in each successive age cohort from the 20s through the 90s. The magnitude of this change was substantial, with the oldest cohort having mental health scores 1 SD higher than those of the youngest cohort, in sharp contrast to the 1.5 and 2 SD differences (suggesting declines) in physical and cognitive function, respectively. The age effects on the composite mental health measure did not conform to a “U-shaped” curve; rather, a linear increase from the 20s to the 90s provided the best fitting model.

Study strengths include the structured multicohort design, subject selection using random digit dialing, oversampling of older adults (due to increased risk of dropouts), and use of published, validated measures of various constructs simultaneously. However, our study also has limitations. First, data were cross-sectional; therefore, we cannot make longitudinal or causal inferences about changes in these domains for individual participants. The apparent changes might reflect birth cohort differences (eg, baby boomers have a higher incidence of mental illnesses including depression, anxiety disorders, and substance use disorders compared to people born before World War II)<sup>46</sup> and/or survivor bias (ie, less healthy younger adults are less likely to survive into old age). However, the negative trajectory of physical and cognitive health across age groups with expected magnitudes of decline argues against the older participants being *super-healthy*. While longitudinal studies to examine aging trajectories require many resources, are expensive, and require more than a decade of follow-up, employment of innovative study designs, such as the multicohort longitudinal design that we used, can make such research more feasible. Second, self-report measures of physical and mental health could be less accurate than objective measures. However, investigations have shown that self-reports of subjective functioning correlate significantly with objective measures of health (eg, mortality)<sup>47</sup> and of local community wellness,<sup>48</sup> suggesting that subjective measures do have objective value. Third, representativeness of the sample might have varied across age groups. While we used random digit dialing to reduce bias in convenience sampling, our study protocol required that participants be selected/contacted using a landline. Many younger adults use cell phones as their main form of communication<sup>49</sup>; we might not have achieved a fully representative sample among

younger age groups. However, cell phone numbers do not correspond to the local area code of residence and, thus, do not help ascertain if the participant is located in San Diego County. Similarly, measurement floor or ceiling effects could have impacted the analyses. Furthermore, we did not use a comprehensive neuropsychological evaluation to measure cognition, but rather a telephone-based screening tool (TICS-M). Although dementia was screened out by self-report, our sample quite likely included some individuals with previously undiagnosed dementia.<sup>32</sup> Some of the observed cognitive decline beginning around age 65 years may reflect such pathological cognitive aging. Finally, our sample was based in San Diego, and further studies of this type are needed in other regions to generalize our findings to communities with different population characteristics.

Notwithstanding these limitations, an intriguing question is whether the observed positive mental health in old age reflects favorable brain changes<sup>50</sup> or is the result of diminished risk factors for mental distress. One possibility is that older adults become more adept at coping with stressful changes.<sup>12,13,51–53</sup> Similar to the literature on cognitive reserve in which passive versus active processes converge in maintaining cognitive function,<sup>54</sup> active emotional reserve may enable older adults to counter threats to mental health such as diminishing physical health. Carstensen and colleagues<sup>14</sup> have attributed such changes to socioemotional selectivity in later life, resulting from narrower horizon of life in view of the acknowledgment of mortality. Another possible explanation for greater life satisfaction among older adults is what Linda George described 30 years ago as “lowering [of] aspirations to meet realities.”<sup>55(p7)</sup>

An important explanation for improved mental health in later life is an increase in wisdom with aging, as suggested by several (but not all) studies.<sup>8,13,56</sup> Researchers have reported that, compared to younger adults, older individuals tend to be more skilled at emotional regulation and complex social decision-making and tend to exhibit more positively valenced information processing.<sup>12–16</sup> Others have found that as people age, they experience fewer negative emotions, regulate their emotions more effectively, and show positive biases in their memory.<sup>57</sup> One study<sup>16</sup> reported that older adults used accumulated lifetime experience in decision-making to determine long-term utility rather than immediate gains. In one investigation, Gooding and colleagues<sup>58</sup> reported that older adults were more resilient than younger ones with respect to emotional regulation and problem solving. These behavioral changes parallel functional imaging studies showing diminished responsiveness of the amygdala to negative or stressful images in older compared to younger adults.<sup>59</sup> Such positivity may lead to higher levels of subjective well-being in later life.

Our results are consistent with epidemiologic studies finding that the prevalence of nondementing mental disorders declines incrementally from younger to older age groups.<sup>22–25</sup> Our results suggest that surviving older adults may be more resilient to common physical and social stresses compared to younger counterparts. A similar phenomenon

may underlie the lower prevalence of mental illnesses in older adults, although there are alternative explanations such as underdiagnosis or misdiagnosis of psychiatric conditions in later life.

At the same time, old age is often associated with serious physical, psychological, and social stresses. One must not underestimate the need for high quality mental and physical health care for older adults. Nonetheless, our study raises important questions about traditional conceptualization of models of aging, including biological ones, that focus on functional (physical and cognitive) declines with usual aging or employ longevity as the primary marker of successful aging. These models need revising to account for what may be “normative” improvement in mental health that may accompany population aging. Even among people with serious mental illnesses such as schizophrenia, aging tends to be associated with improvement in mental health.<sup>60,61</sup>

Another notable finding relates to high levels of subjective (perceived) stress along with symptoms of depression and anxiety among those in their twenties and thirties. This

“fountain of youth” is associated with a far worse level of psychological well-being than during any other period of adulthood. There are many pressures rather unique to this life phase—eg, establishing careers, finding life partners, and navigating financial issues.<sup>62</sup> Whereas adolescence has long been an area of social and scientific concern, relatively little attention has been paid to issues that continue or get exacerbated postadolescence. Of note, the rate of mental illness and treatment seeking in younger persons has been increasing sharply,<sup>63</sup> whereas rates of self-reported depressive symptoms in later life have declined.<sup>64</sup>

When treating younger adults, we should consider how to leverage positive mental health factors to augment their psychosocial functioning, whereas in older adults, we should consider how these factors may help with physical and cognitive functioning. There is a critical need for lifespan research combining psychosocial and biological sciences to improve our understanding of the processes that underlie increased well-being in later life and to help develop interventions to promote wellness in all age groups.

**Submitted:** January 13, 2016; accepted May 3, 2016.

**Potential conflicts of interest:** None to declare.

**Funding/support:** Supported, in part, by National Institute of Mental Health grants T32 MH019934, R01 MH099987, R01 MH094151, and K23 MH102420 and by the Sam and Rose Stein Institute for Research on Aging at the University of California, San Diego.

**Role of the sponsor:** The donors to the Stein Institute had no role in the design and conduct of this study.

**Previous presentation:** None.

**Acknowledgments:** We thank Rebecca Daly, AA, from the Department of Psychiatry and the Sam and Rose Stein Institute for Research on Aging at University of California, San Diego, for invaluable contributions in database management. She has no conflicts of interest to report.

**Supplementary material:** See accompanying pages.

## REFERENCES

- Executive Summary: A Survey About Mental Health and Suicide in the United States. American Foundation for Suicide Prevention Web site. <http://afsp.org/executive-summary-survey-mental-health-suicide-united-states/>. Accessed September 3, 2015.
- Mental disorders. WHO Web site. <http://www.who.int/mediacentre/factsheets/fs396/en/>. Accessed August 21, 2015.
- Moussavi S, Chatterji S, Verdes E, et al. Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet*. 2007;370(9590):851–858.
- Steptoe A, Deaton A, Stone AA. Subjective wellbeing, health, and ageing. *Lancet*. 2015;385(9968):640–648.
- Jeste DV, Depp CA, Vahia IV. Successful cognitive and emotional aging. *World Psychiatry*. 2010;9(2):78–84.
- Blanchflower DG, Oswald AJ. Is well-being U-shaped over the life cycle? *Soc Sci Med*. 2008;66(8):1733–1749.
- Stone AA, Schwartz JE, Broderick JE, et al. A snapshot of the age distribution of psychological well-being in the United States. *Proc Natl Acad Sci U S A*. 2010;107(22):9985–9990.
- Jeste DV, Oswald AJ. Individual and societal wisdom: explaining the paradox of human aging and high well-being. *Psychiatry*. 2014;77(4):317–330.
- López Ulloa BF, Möller V, Sousa-Poza A. How does subjective well-being evolve with age? a literature review. *J Population Ageing*. 2013;6(3):227–246.
- Charles ST, Reynolds CA, Gatz M. Age-related differences and change in positive and negative affect over 23 years. *J Pers Soc Psychol*. 2001;80(1):136–151.
- Mroczek DK, Kolarz CM. The effect of age on positive and negative affect: a developmental perspective on happiness. *J Pers Soc Psychol*. 1998;75(5):1333–1349.
- Blazer DG. Successful aging. *Am J Geriatr Psychiatry*. 2006;14(1):2–5.
- Vaillant GE. *Aging Well: Surprising Guideposts to a Happier Life From the Landmark Harvard Study of Adult Development*. Boston, MA: Little, Brown & Company; 2002.
- Carstensen LL, Mikels JA, Mather M. Aging and the intersection of cognition, motivation, and emotion. In: Birren JE, Schaie KW, Abeles RP, et al. *Handbook of the Psychology of Aging*. Burlington, MA: Elsevier; 2006:343–362.
- Grossmann I, Na J, Varnum MEW, et al. Reasoning about social conflicts improves into old age. *Proc Natl Acad Sci U S A*. 2010;107(16):7246–7250.
- Worthy DA, Gorlick MA, Pacheco JL, et al. With age comes wisdom: decision making in younger and older adults. *Psychol Sci*. 2011;22(11):1375–1380.
- Springer KW, Pudrovskaya T, Hauser RM. Does psychological well-being change with age? longitudinal tests of age variations and further exploration of the multidimensionality of Ryff's Model of Psychological Well-Being. *Soc Sci Res*. 2011;40(1):392–398.
- Wild S, Roglic G, Green A, et al. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004;27(5):1047–1053.
- Ho KK, Pinsky JL, Kannel WB, et al. The epidemiology of heart failure: the Framingham Study. *J Am Coll Cardiol*. 1993;22(6 suppl A):6A–13A.
- Craik FIM, Salthouse TA. *The Handbook of Aging and Cognition*. Mahwah, NJ: Lawrence Erlbaum Associates; 2000.
- Andrews FM, Withey SB. *Social Indicators of Well-Being: Americans' Perceptions of Life Quality*. New York, NY: Plenum Press; 1976.
- Hasin DS, Goodwin RD, Stinson FS, et al. Epidemiology of major depressive disorder: results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Arch Gen Psychiatry*. 2005;62(10):1097–1106.
- de Graaf R, ten Have M, van Gool C, et al. Prevalence of mental disorders and trends from 1996 to 2009: results from the Netherlands Mental Health Survey and Incidence Study-2. *Soc Psychiatry Psychiatr Epidemiol*. 2012;47(2):203–213.
- Regier DA, Farmer ME, Rae DS, et al. One-month prevalence of mental disorders in the United States and sociodemographic characteristics: the Epidemiologic Catchment Area study. *Acta Psychiatr Scand*. 1993;88(1):35–47.
- Kessler RC, McGonagle KA, Zhao S, et al. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry*. 1994;51(1):8–19.
- Palmer BW, Jeste DV, Sheikh JI. Anxiety disorders in the elderly: DSM-IV and other barriers to diagnosis and treatment. *J Affect Disord*. 1997;46(3):183–190.
- Jeste DV, Savla GN, Thompson WK, et al. Association between older age and more successful aging: critical role of resilience and depression. *Am J Psychiatry*. 2013;170(2):188–196.
- Martin AS, Palmer BW, Rock D, et al. Associations of self-perceived successful aging in young-old versus old-old adults. *Int Psychogeriatr*. 2015;27(4):601–609.
- Ware JE, Kosinski M, Keller S. *SF-36 Physical and Mental Health Summary Scales: A User's Manual*. 1994.
- Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36), I: conceptual framework and item selection. *Med Care*. 1992;30(6):473–483.
- de Jager CA, Budge MM, Clarke R. Utility of TICS-M for the assessment of cognitive function in older adults. *Int J Geriatr Psychiatry*.

- 2003;18(4):318–324.
32. Zlatar ZZ, Moore RC, Palmer BW, et al. Cognitive complaints correlate with depression rather than concurrent objective cognitive impairment in the successful aging evaluation baseline sample. *J Geriatr Psychiatry Neurol*. 2014;27(3):181–187.
33. Fowler JH, Christakis NA. Dynamic spread of happiness in a large social network: longitudinal analysis over 20 years in the Framingham Heart Study. *BMJ*. 2008;337:a2338.
34. Diener E, Emmons RA, Larsen RJ, et al. The Satisfaction With Life Scale. *J Pers Assess*. 1985;49(1):71–75.
35. Derogatis LR. *Brief Symptom Inventory: Administration, Scoring, and Procedures Manual*. 4th ed. Minneapolis, MN: National Computer Systems; 1993.
36. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606–613.
37. Cohen S, Williamson G. Perceived stress in a probability sample of the United States. In: Spacapan S, Oskamp S, eds. *The Social Psychology of Health*. Newbury Park, CA: SAGE Publications; 1988:31–67.
38. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983;24(4):385–396.
39. Cohen J, Cohen P, West SG, et al. *Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers; 2003.
40. Serlin RC, Levin JR. Teaching how to derive directly interpretable coding schemes for multiple regression analysis. *J Educ Behav Stat*. 1985;10(3):223–238.
41. R Development Core Team. R: A Language and Environment for Statistical Computing [computer program]. Vienna, Austria: R Foundation for Statistical Computing; 2014.
42. Glenn N. Is the apparent U-shape of well-being over the life course a result of inappropriate use of control variables? a commentary on Blanchflower and Oswald (66: 8, 2008, 1733–1749). *Soc Sci Med*. 2009;69(4):481–485, discussion 486–488.
43. Nair KS. Aging muscle. *Am J Clin Nutr*. 2005;81(5):953–963.
44. Lunney JR, Lynn J, Foley DJ, et al. Patterns of functional decline at the end of life. *JAMA*. 2003;289(18):2387–2392.
45. Fleg JL, Morrell CH, Bos AG, et al. Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation*. 2005;112(5):674–682.
46. Jeste DV, Alexopoulos GS, Bartels SJ, et al. Consensus statement on the upcoming crisis in geriatric mental health: research agenda for the next 2 decades. *Arch Gen Psychiatry*. 1999;56(9):848–853.
47. DeSalvo KB, Bloser N, Reynolds K, et al. Mortality prediction with a single general self-rated health question: a meta-analysis. *J Gen Intern Med*. 2006;21(3):267–275.
48. Oswald AJ, Wu S. Objective confirmation of subjective measures of human well-being: evidence from the USA. *Science*. 2010;327(5965):576–579.
49. Lee S, Brick JM, Brown ER, et al. Growing cell-phone population and noncoverage bias in traditional random digit dial telephone health surveys. *Health Serv Res*. 2010;45(4):1121–1139.
50. Charles ST, Carstensen LL. Social and emotional aging. *Annu Rev Psychol*. 2010;61(1):383–409.
51. Jeste DV, Palmer BW. *Positive Psychiatry—a Clinical Handbook*. Washington, DC: American Psychiatric Publishing; 2015.
52. Jeste DV, Palmer BW, Rettew DC, et al. Positive psychiatry: its time has come. *J Clin Psychiatry*. 2015;76(6):675–683.
53. Reynolds CF 3rd. Preventing depression in old age: it's time. *Am J Geriatr Psychiatry*. 2008;16(6):433–434.
54. Stern Y. Cognitive reserve. *Neuropsychologia*. 2009;47(10):2015–2028.
55. George LK. Life satisfaction in later life. *Generations: Journal of the American Society on Aging*. 1986.
56. Mickler C, Staudinger UM. Personal wisdom: validation and age-related differences of a performance measure. *Psychol Aging*. 2008;23(4):787–799.
57. Reed AE, Carstensen LL. The theory behind the age-related positivity effect. *Front Psychol*. 2012;3:339.
58. Gooding PA, Hurst A, Johnson J, et al. Psychological resilience in young and older adults. *Int J Geriatr Psychiatry*. 2012;27(3):262–270.
59. Mather M, Canli T, English T, et al. Amygdala responses to emotionally valenced stimuli in older and younger adults. *Psychol Sci*. 2004;15(4):259–263.
60. Jeste DV, Wolkowitz OM, Palmer BW. Divergent trajectories of physical, cognitive, and psychosocial aging in schizophrenia. *Schizophr Bull*. 2011;37(3):451–455.
61. Folsom DP, Depp C, Palmer BW, et al. Physical and mental health-related quality of life among older people with schizophrenia. *Schizophr Res*. 2009;108(1–3):207–213.
62. Schwartz B. *The Paradox of Choice*. New York, NY: Harper Collins; 2009.
63. Collishaw S. Annual research review: secular trends in child and adolescent mental health. *J Child Psychol Psychiatry*. 2015;56(3):370–393.
64. Zivin K, Pirraglia PA, McCammon RJ, et al. Trends in depressive symptom burden among older adults in the United States from 1998 to 2008. *J Gen Intern Med*. 2013;28(12):1611–1619.

---

Supplementary material follows this article.

---



## **Supplementary Material**

**Article Title:** Paradoxical Trend for Improvement in Mental Health With Aging: A Community-Based Study of 1,546 Adults Aged 21–100 Years

**Author(s):** Michael L. Thomas, PhD; Christopher N. Kaufmann, PhD, MHS; Barton W. Palmer, PhD; Colin A. Depp, PhD; Averria Sirkin Martin, PhD; Danielle K. Glorioso, MSW; Wesley K. Thompson, PhD; and Dilip V. Jeste, MD

**DOI Number:** 10.4088/JCP.16m10671

### **List of Supplementary Material for the article**


1. [eFigure 1](#) CONSORT Diagram of Participants Enrolled in the SAGE Study
2. [eTable 1](#) Demographic and Clinical Characteristics (N = 1,546)
3. [eFigure 2](#) Scatterplots for Each Domain
4. [Supplemental Text](#)

### **Disclaimer**

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.



**eFigure 1. CONSORT Diagram of Participants Enrolled in the SAGE Study**




a. Recruitment full, no answer, answering machine on repeated calls, etc.

b. Disconnected number, didn't speak English, deceased, hard of hearing, etc.

**eTable 1. Demographic and Clinical Characteristics (N = 1,546)**

Age mean ( <i>SD</i> ) in years	66 (21)
Male <i>N</i> (%)	782 (51%)
Ethnicity/Race <i>N</i> (%)	
White	1,175 (76%)
Hispanic/Latino	217 (14%)
Asian-American	108 (7%)
African American	23 (1%)
Other	23 (1%)
Education <i>N</i> (%)	
12 years or less	302 (20%)
13-16 years	917 (60%)
17 years or more	322 (21%)

**eFigure 2. Scatterplots for Each Domain**



### **Supplemental Text**

Participants were asked questions pertaining to anxiety and depression while on the phone with an interviewer during the initial screening and telephone interview. Interviewers were carefully trained to immediately inform the investigators if a participant expressed suicidal ideation, thoughts of self-harm, or other concerns warranting clinical attention, at which point one of the licensed clinician co-investigators would follow-up with the participant to properly assess and guide any warranted follow-up steps to ensure participant safety.

Although the overall effects of age were generally moderate to large, it should be emphasized that some of the specific terms were not. Specifically, the quadratic term in the physical health model as well as the cubic term in the cognitive function model were both statistically significant but the effects were small.