

Supplementary Material

Article Title: Prevalence and Correlates of Obsessive-Compulsive Symptoms in Individuals With

Schizophrenia, Schizoaffective Disorder, or Bipolar Disorder

Authors: Deborah Ahn-Robbins, MSc; Nina H. Grootendorst-van Mil, MD, PhD; Chin-Kuo Chang, PhD;

David Chandran, PhD; Hitesh Shetty, PhD; Jyoti Sanyal, PhD;

James H. MacCabe, FRCPsych, PhD; Hannah Cohen, PhD; Robert Stewart, MD; Frederike Schirmbeck, PhD; Lieuwe de Haan, MD, PhD; and Richard D. Hayes, PhD

DOI Number: 10.4088/JCP.21m14010

List of Supplementary Material for the article

1. Table 1 Associations Between Mental/Physical Health Symptoms and OCS (Including OCD)

Status Among SLaM Service Users With Schizophrenia, Schizoaffective Disorder or Bipolar Disorder in Southeast London by Univariable and Multivariable Logistic

Regression

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Table 1. Associations between mental / physical health symptoms and OCS (including OCD) status among SLaM service users with schizophrenia, schizoaffective disorder or bipolar disorder in southeast London by univariable and multivariable logistic regression

Odds Ratio (95% CI)

Status of obsessive	Overactive aggressive behaviour ^a		Non-accidental self-injury b		Cognitive problems ^c	
compulsive symptom	Crude	Adjusted^	Crude	Adjusted^	Crude	Adjusted^
OCS (including OCD)						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	1.18 (1.11, 1.26)	1.17 (1.9, 1.25)	1.18 (1.08, 1.29)	1.04 (0.95, 1.14)	1.14 (1.07, 1.23)	1.20 (1.12, 1.29)
(continue)						
	Odds Ratio (95% CI)					
Status of obsessive	Physical illness or disability ^d		Hallucinations and delusions ^e		Depressed mood f	
compulsive symptom	Crude	Adjusted^	Crude	Adjusted^	Crude	Adjusted^
OCS (including OCD)						
No	Ref	Ref	Ref	Ref	Ref	Ref
Yes	0.98 (0.92, 1.05)	1.17 (1.09, 1.27)	1.11 (1.04, 1.19)	1.09 (1.02, 1.18)	1.09 (1.02, 1.17)	1.05 (0.98, 1.12)

[^] Adjusting age, sex, ethnicity, relationship, deprivation score, psychiatric diagnosis (schizophrenia, schizoaffective disorder, or bipolar disorder), comorbidity of alcohol use disorder, and comorbidity of opioid use disorder

^a N=18,035; ^b N=18,007; ^c N=17,983; ^d N=17,995; ^e N=17,955; ^f N=17,982