



# THE JOURNAL OF CLINICAL PSYCHIATRY

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

## Supplementary Material

**Title:** Changes in Alcohol-Related Behaviors and Quality of Life During the COVID-19 Pandemic: Impact of Alcohol Use Disorder Diagnosis and Treatment History

**Authors:** Jeremy W. Luk, PhD; Bethany L. Stangl, PhD; Tommy Gunawan, PhD; Melanie L. Schwandt, PhD; James K. Morris, BS; Jared C. Axelowitz, BS; Sumedha Chawla, MBBS, MPH; Beth A. Lee, MSN; Megan Carraco, MS; Sheila Walsh, MA, RN-PMHN-BC; Christian McDuffie, BA; Khushbu Agarwal, PhD; Paule V. Joseph, PhD, MS, FNP-BC, CTN-B, FAAN; Reza Momenan, PhD; David Goldman, MD; Nancy Diazgranados, MD, MS; and Vijay A. Ramchandani, PhD

**DOI Number:** 10.4088/JCP.22br14462

### List of Supplementary Material for the Brief Report

1. [Table 1](#) Effects of Time on Alcohol-Related and Quality of Life Outcomes by AUD Treatment Status

### Disclaimer

This Supplementary Material has been provided by the authors as an enhancement to the published brief report. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

It is illegal to post this copyrighted PDF on any website. ♦ © 2023 Copyright Physicians Postgraduate Press, Inc.

© Copyright 2022 Physicians Postgraduate Press, Inc.

Supplementary Table 1. Effects of Time on Alcohol-Related and Quality of Life Outcomes by AUD-Treatment Status

	A. Problematic Drinking		B. Alcohol Craving		C. Physical Quality of Life	
	<i>b</i> (95% CIs)	<i>p</i>	<i>b</i> (95% CIs)	<i>p</i>	<i>b</i> (95% CIs)	<i>p</i>
Healthy Control						
T2 vs T1	0.13 (-0.89, 1.15)	0.805	<b>1.72 (0.69, 2.76)</b>	<b>0.001</b>	<b>-5.13 (-7.47, -2.79)</b>	<b>&lt;0.001</b>
T3 vs T1	0.07 (-0.95, 1.09)	0.895	<b>2.69 (1.65, 3.73)</b>	<b>&lt;0.001</b>	<b>-12.51 (-14.85, -10.17)</b>	<b>&lt;0.001</b>
T3 vs T2	-0.06 (-1.08, 0.96)	0.907	0.96 (-0.06, 1.98)	0.065	<b>-7.38 (-9.67, -5.09)</b>	<b>&lt;0.001</b>
Non-Treatment AUD						
T2 vs T1	<b>-2.84 (-4.88, -0.81)</b>	<b>0.006</b>	<b>-2.68 (-4.62, -0.75)</b>	<b>0.007</b>	-2.54 (-7.00, 1.91)	0.263
T3 vs T1	<b>-2.56 (-4.58, -0.53)</b>	<b>0.013</b>	-0.17 (-2.11, 1.76)	0.860	<b>-14.89 (-19.35, -10.44)</b>	<b>&lt;0.001</b>
T3 vs T2	0.29 (-1.74, 2.32)	0.781	<b>2.51 (0.59, 4.43)</b>	<b>0.011</b>	<b>-12.35 (-16.74, -7.96)</b>	<b>&lt;0.001</b>
Treatment AUD						
T2 vs T1	<b>-11.99 (-13.43, -10.55)</b>	<b>&lt;0.001</b>	<b>5.51 (3.99, 7.04)</b>	<b>&lt;0.001</b>	1.97 (-1.76, 5.70)	0.300
T3 vs T1	<b>-13.76 (-15.20, -12.32)</b>	<b>&lt;0.001</b>	<b>4.82 (3.29, 6.34)</b>	<b>&lt;0.001</b>	-1.08 (-4.80, 2.65)	0.571
T3 vs T2	<b>-1.77 (-3.19, -0.34)</b>	<b>0.015</b>	-0.69 (-2.16, 0.77)	0.353	-3.05 (-6.74, 0.64)	0.106
	D. Psychological Quality of Life		E. Social Quality of Life		F. Environment Quality of Life	
	<i>b</i> (95% CIs)	<i>p</i>	<i>b</i> (95% CIs)	<i>p</i>	<i>b</i> (95% CIs)	<i>p</i>
Healthy Control						
T2 vs T1	<b>-7.87 (-10.55, -5.18)</b>	<b>&lt;0.001</b>	<b>-8.90 (-12.28, -5.52)</b>	<b>&lt;0.001</b>	<b>-4.05 (-6.52, -1.57)</b>	<b>0.001</b>
T3 vs T1	<b>-15.96 (-18.65, -13.28)</b>	<b>&lt;0.001</b>	<b>-12.53 (-15.90, -9.15)</b>	<b>&lt;0.001</b>	<b>-11.27 (-13.74, -8.80)</b>	<b>&lt;0.001</b>
T3 vs T2	<b>-8.10 (-10.73, -5.47)</b>	<b>&lt;0.001</b>	<b>-3.63 (-6.93, -0.32)</b>	<b>0.032</b>	<b>-7.22 (-9.64, -4.80)</b>	<b>&lt;0.001</b>
Non-Treatment AUD						
T2 vs T1	<b>-5.72 (-10.84, -0.60)</b>	<b>0.029</b>	-4.61 (-11.05, 1.83)	0.161	2.19 (-2.53, 6.90)	0.363
T3 vs T1	<b>-17.52 (-22.64, -12.41)</b>	<b>&lt;0.001</b>	<b>-13.81 (-20.26, -7.37)</b>	<b>&lt;0.001</b>	<b>-5.76 (-10.47, -1.04)</b>	<b>0.017</b>
T3 vs T2	<b>-11.81 (-16.85, -6.76)</b>	<b>&lt;0.001</b>	<b>-9.20 (-15.55, -2.85)</b>	<b>0.005</b>	<b>-7.94 (-12.59, -3.30)</b>	<b>&lt;0.001</b>
Treatment AUD						
T2 vs T1	<b>5.51 (1.23, 9.79)</b>	<b>0.012</b>	1.62 (-3.77, 7.01)	0.556	1.64 (-2.31, 5.58)	0.416
T3 vs T1	4.10 (-0.18, 8.38)	0.061	3.33 (-2.06, 8.72)	0.225	-1.40 (-5.34, 2.55)	0.487
T3 vs T2	-1.41 (-5.65, 2.83)	0.515	1.72 (-3.62, 7.05)	0.529	-3.03 (-6.94, 0.87)	0.128

Multilevel models adjusted for age, sex, race, cohort, and enrollment phase. Regression coefficients were unstandardized. The ranges of possible scores were from 0 to 40 for the Alcohol Use Disorders Identification Test, 0 to 30 for Penn Alcohol Craving Scale, and 0 to 100 for each of the four World Health Organization QoL - BREF domains.