



Supplementary Material

Article Title: Bad Dreams and Nightmares Preceding Suicidal Behaviors

Author(s): Pierre A. Geoffroy, MD, PhD; Rodolphe Borand, MD; Marine Ambar Akkaoui, MD, MSc; Séverine Yung, MD; Yasmine Atoui, MD; Emeline Fontenoy, MD; Julia Maruani, MD, MSc; and Michel Lejoyeux, MD, PhD

DOI Number: 10.4088/JCP.22m14448

List of Supplementary Material for the article

1. [Table 1](#) Dreams Content and Suicidal Phenotypes of Patient Admitted in a University Psychiatric Post-Emergency Room Unit for Suicidal Crisis
2. [Figure 1](#) Altered Dreams in Patients With (A) Suicidal Crisis (Thoughts and Attempts) and (B) Suicide Attempt


Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Table 1. Dreams content and suicidal phenotypes of patient admitted in a University psychiatric post-emergency room unit for suicidal crisis.

Total sample n=40	Suicidal ideation n=40	Suicide attempts n=29	Recurrence of suicide attempts n=18
No altered dreams n=8	8 (20%)	6 (20.7%)	4 (22.2%)
Altered dreams n=32	32 (80%)	23 (79.3%)	14 (77.8%)
Bad dreams n=27	27 (67.5%)	20 (69.0%)	11 (61.1%)
Nightmares n=21	21 (52.5%)	14 (48.3%)	9 (50%)
Suicidal scenarii during dreams n= 9	9 (22.5%)	6 (20.7%)	4 (22.2%)

Supplementary Figure 1
(A) suicidal crisis (thoughts and attempts)



Supplementary Figure 1
(B) suicide attempt

