

Supplementary Material

Article Title: Sleep Complaints Among Adults With Major Depressive Episode Are Associated With

Increased Risk of Incident Psychiatric Disorders: Results From a Population-Based 3-Year

Prospective Study

Authors: Bénédicte Barbotin, MD; Nicolas Hoertel, MD, MPH, PhD; Mark Olfson, MD, MPH;

Carlos Blanco, MD, PhD; Marina Sanchez-Rico, MPH; Michel Lejoyeux, MD, PhD;

Frédéric Limosin, MD, PhD; and Pierre A. Geoffroy, MD, PhD

DOI Number: 10.4088/JCP.21m14236

List of Supplementary Material for the article

1. Table 1 Single Factor Confirmatory Factor Analysis (CFA) of Lifetime Sleep Complaints (ie,

Trouble Falling Asleep, Early Wake-Up, Hypersomnia) Among Participants With a Major Depressive Episode in The Year Prior to Wave 1 of the National Epidemiologic Survey on

Alcohol and Related Conditions (NESARC) (N=2864)

2. Table 2 Bifactor Model of the Structure Distress-Fear-Externalizing Underlying Past-Year Axis I

Disorders and Antisocial Personality Disorder Among Participants With a Major

Depressive Episode in the Year Prior to Wave 1 of the National Epidemiologic Survey on

Alcohol and Related Conditions (NESARC) (N=2864)

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Table 1. Single factor confirmatory factor analysis (CFA) of lifetime sleep complaints (i.e., trouble falling asleep, early wake-up, hypersomnia) among participants with a major depressive episode in the year prior to Wave 1 of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC) (N=2864).

CFI	0.99			
TLI	0.99			
RMSEA	0.01			
	Sleep complaints			
Trouble falling asleep	0.71**			
Early wake-up	0.83**			
Hypersomnia	-0.52**			

Abbreviations: CFI, Comparative Fit Index; TLI, Tucker-Lewis Index; RMSEA, Root Mean Square Error of Approximation. ** p-value<0.01.

Supplementary Table 2. Bifactor model of the structure distress-fear-externalizing underlying past-year Axis I disorders and antisocial personality disorder among participants with a major depressive episode in the year prior to Wave 1 of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC) (N=2864).

CFI	0.99					
TLI	0.99 0.01					
RMSEA						
Psychiatric disorders	General Psychopathology factor	Distress	Fear	Externalizing		
Dysthymia	0.50**	0.71**				
General anxiety disorder	0.34**	0.26*				
Panic disorder	0.49**		0.68**			
Social anxiety disorder	0.86**		-0.20			
Specific phobia	0.47**		-0.04			
Alcohol use disorder	0.20*			0.69**		
Nicotine dependence	0.14			0.37**		
Cannabis use disorder	0.48**			0.58**		
Drug use disorder	0.43**			0.47**		
Factor loading of the factor Internalizing		0.71**	0.60*			
Factor correlation						
General Psychopathology factor	1.00					
Distress	0.00	1.00				
Fear	0.00	0.00	1.00			
Externalizing	0.00	0.00	0.00	1.00		

Abbreviations: CFI, Comparative Fit Index; TLI, Tucker-Lewis Index; RMSEA, Root Mean Square Error of Approximation.

^{*} p-value<0.05; ** p-value<0.01.