



THE JOURNAL OF
CLINICAL PSYCHIATRY

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

Supplementary Material

Article Title: Impact of Tardive Dyskinesia on Physical, Psychological, Social, and Professional Domains of Patient Lives: A Survey of Patients in the United States

Author(s): Rakesh Jain, MD, MPH; Rajeev Ayyagari, PhD; Debbie Goldschmidt, PhD; Mo Zhou, PhD; Stacy Finkbeiner, PhD; and Sam Leo, PharmD

DOI Number: <https://doi.org/10.4088/JCP.22m14694>

List of Supplementary Material for the article


1. [Figure 1](#) Patient Impact Overall and by TD Symptom Severity or Underlying Condition
2. [Figure 2](#) Social Impact on the Patient of Reactions From Acquaintances
3. [Table 1](#) Patient-reported Coping Strategies for TD

Disclaimer


This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Figure 1. Patient Impact Overall and by TD Symptom Severity or Underlying Condition

A. By TD symptom severity



B. By underlying condition




Abbreviations: TD=tardive dyskinesia.

Supplementary Figure 2. Social Impact on the Patient of Reactions From Acquaintances

How often do people that you know (such as family, friends, or coworkers) “_____”?

1. Stare or look at you because of TD
2. Ask what is wrong or why you are moving
3. Tell you to stop moving
4. Make jokes at your expense because of TD
5. Tell you that you are different than how you used to be because of TD
6. Grab or touch you to try to stop you from moving
7. Tell you that other people are looking at you because of TD



Abbreviation: TD=tardive dyskinesia.

Supplementary Table 1. Patient-reported Coping Strategies for TD

Strategies to cope with TD movements^a	Total N=269 n (%)
Slow down when speaking	151 (56.1)
Massage, stretch, or shake out body parts that are cramping or moving	133 (49.4)
Take prescription medications to relax muscles	131 (48.7)
Take sleeping pills or sleeping aids	125 (46.5)
Cover their hands	118 (43.9)
Get up and walk around when movements start	116 (43.1)
Meditate or do breathing exercises like deep or slow breathing	112 (41.6)
Suck on hard candy or chew gum	95 (35.3)
Cover their face	86 (32.0)
Postpone activity until extra movements subside	84 (31.2)

^aRespondents could select more than one option; results are not mutually exclusive.
Abbreviation: TD=tardive dyskinesia.