

Supplementary Material

Article Title: A Comparison of Depressive Symptom Self-Reported Measures in the Texas Youth Depression and Suicide Research Network (TX-YDSRN)

Authors: Karabi Nandy, PhD; A. John Rush, MD; Thomas Carmody, PhD; Beth D. Kennard, MD; Graham J. Emslie, MD; Holli Slater, PhD; Taryn L. Mayes, MS; Melissa DeFilippis, MD; Cynthia Garza, PsyD; Eric A. Storch, PhD; Sarah M. Wakefield, MD; and Madhukar H. Trivedi, MD

DOI Number: 10.4088/JCP.23m14861

LIST OF SUPPLEMENTARY MATERIAL FOR THE ARTICLE

1. [Table 1](#) List of Items in IDS-SR, QIDS-SR₁₆, PHQ-A, VQIDS-SR₅
2. [Figure 1](#) Frequencies of QIDS-SR₁₆ and PHQ-A Ratings by Domain (n=795)
3. [Figure 2](#) Scatterplots of Change Scores of Pairs of PHQ-A, QIDS-SR₁₆ and VQIDS-SR₅ Total Scores (n=682)

DISCLAIMER

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Three Scales for Measuring Adolescent Depression

SUPPLEMENTARY TABLES AND FIGURES


Supplementary Table 1. List of items in IDS-SR, QIDS-SR₁₆, PHQ-A, VQIDS-SR₅ (all items are scores 0-3)

30-item IDS-SR	16-item QIDS-SR ₁₆	PHQ-A (matching 9 domains of PHQ-A to the 9 domains in QIDS-SR ₁₆)	5-item VQIDS-SR ₅
1. Sleep onset insomnia	√	Item 3: Trouble falling or staying asleep, or sleeping too much	
2. Mid-nocturnal insomnia	√	Item 3	
3. Early morning insomnia	√	Item 3	
4. Hypersomnia	√	Item 3	
5. Mood (sad)	√	Item 2: Feeling down, depressed, or hopeless	√
6. Mood (irritable)			
7. Feeling anxious or tense			
8. Response of Mood to Good or Desired Events			
9. Mood in relation to time of day			
10. Quality of mood			
11. Appetite (decreased)	√	Item 5: Poor appetite or overeating	
12. Appetite (increased)	√	Item 5	
13. Weight (decrease) within the last two weeks	√	Item 5	
14. Weight (increase) within the last two weeks	√	Item 5	
15. Concentration/decision making	√	Item 7: Trouble concentrating	
16. Outlook (self)	√	Item 6: Feeling bad about yourself – or that you are a failure	√
17. View of future			
18. Suicidal ideation (Thoughts of Death or Suicide)	√	Item 9: Thoughts that you would be better off dead or of hurting yourself	
19. Involvement (General Interest)	√	Item 1: Little interest or loss of pleasure	√
20. Energy/fatigability	√	Item 4: Feeling tired or having little energy	√
21. Capacity for Pleasure or Enjoyment (excluding sex)			
22. Interest in Sex			
23. Psychomotor slowing (Feeling slowed down)	√	Item 8: Moving or speaking so slowly that other people could notice	√
24. Psychomotor agitation (Feeling restless)	√	Item 8	
25. Aches and pains			
26. Other bodily symptoms			
27. Panic/Phobic symptoms			
28. Constipation/diarrhea			
29. Interpersonal Sensitivity			
30. Leadon Paralysis/Physical Energy			
IDS total score = sum of items (5-10, 15-22, 25 + 26-30) + Q1 + Q2 + Q3; where Q1=Max of 4 sleep items (1 to 4) Q2=Max of 4 appetite/weight items (11 to 14)	QIDS total score = item 5 + item 15 + item 16 + item 18 + item 19 + item 20 + Q1 + Q2 + Q3; where Q1=Max of 4 sleep items (1 to 4) Q2=Max of 4 appetite/weight items (11 to 14)	PHQ Total Score = sum of items 1-9	VQIDS total score = item 5 + item 16 + item 19 + item 20 + item 23


Three Scales for Measuring Adolescent Depression

Q3=Max of 2 psychomotor items (23 and 24)	Q3=Max of psychomotor items (23 and 24)		
---	---	--	--

Supplementary Figure 1. Frequencies of QIDS-SR₁₆ and PHQ-A Ratings by Domain (n=795)



Three Scales for Measuring Adolescent Depression



Supplementary Figure 2: Scatterplots of Change Scores of Pairs of PHQ-A, QIDS-SR₁₆ and VQIDS-SR₅ total scores (n=682)

Three Scales for Measuring Adolescent Depression

