



THE JOURNAL OF CLINICAL PSYCHIATRY

THE OFFICIAL JOURNAL OF THE AMERICAN SOCIETY OF CLINICAL PSYCHOPHARMACOLOGY

Supplementary Material

Article Title: Relapse After Antipsychotic Discontinuation in Schizophrenia as a Withdrawal Phenomenon vs Illness Recurrence: A Post Hoc Analysis of a Randomized Placebo-Controlled Study

Author(s): Robin Emsley, DMed, DSc; Isaac Nuamah, PhD; Srihari Gopal, MD, MHS; David Hough, MD; and W. Wolfgang Fleischhacker, MD

DOI Number: <https://doi.org/10.4088/JCP.17m11874>

List of Supplementary Material for the article

1. [Table 1](#) Clinical Details at Endpoint for the Placebo Patients Who Relapsed Early (within 3 months of randomization) Versus Those Who Relapsed Later (> 3 months after randomization)

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplemental Table: Clinical details at endpoint for the placebo patients who relapsed early (within 3 months of randomization) versus those who relapsed later (> 3 months after randomization)

Characteristic	Early Relapsers (n = 62)	Late Relapsers (n = 35)	P-value
Total PANSS Score	79.4 ± 17.08	77.0 ± 17.48	0.519
Positive Symptoms	23.1 ± 5.55	22.8 ± 5.76	0.806
Negative Symptoms	18.5 ± 5.94	17.1 ± 5.16	0.237
Disorganized Thoughts	18.1 ± 4.76	18.1 ± 4.32	0.981
Uncontrolled Hostility/ Excitement	9.4 ± 3.29	9.4 ± 3.86	0.997
Anxiety/Depression	10.4 ± 2.81	9.7 ± 3.33	0.338
Pulse Rate (bpm) ^a	75.4 ± 10.95	76.9 ± 12.36	0.540
Systolic Blood Pressure (mmHg) ^a	119.1 ± 10.43	124.9 ± 11.32	0.013
Diastolic Blood Pressure (mmHg) ^a	75.0 ± 7.87	78.1 ± 8.66	0.085
Total AIMS	0.39 ± 1.11	0.83 ± 2.37	0.227
Last Antipsychotic Dose (mg)	82.7 ± 25.84	85.0 ± 24.40	0.663
Baseline Prolactin (ng/ml)	47.5 ± 34.91	48.3 ± 44.96	0.915
Prolactin at Endpoint (ng/ml)	43.4 ± 32.67	28.7 ± 36.30	0.050

^aThese are Supine measures. Similar conclusions obtained for standing measures

PANSS, Positive and Negative Syndrome Scale; AIMS, Abnormal Involuntary Movement Scale (to measure dyskinesia)