



Supplementary Material

Article Title: Symptoms of Apathy Independently Predict Incident Frailty and Disability in Community-Dwelling Older Adults

Author(s): Emmeline Ayers, MPH^a; Miriam Shapiro, PhD^a; Roee Holtzer, PhD^{a,b}; Nir Barzilai, MD^{c,d}; Sofiya Milman, MD^c; and Joe Verghese, MBBS, MS^{a,c,*}

DOI Number: <https://doi.org/10.4088/JCP.15m10113>

List of Supplementary Material for the article

1. [eTable 1](#) Results of Cox analysis examining 'dose-dependent' effect of apathy on study outcomes in pooled sample

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary eTable 1. Results of Cox analysis examining 'dose-dependent' effect of apathy on study outcomes

in pooled sample

	Slow Gait	Frailty	Disability
Apathy symptoms endorsed*	N = 822	N = 805	N = 899
1 apathy symptom	1.57 (.98-2.52) .062	1.54 (1.00-2.38) .051	2.52 (1.06=5.99) .036
2 apathy symptoms	2.40 (1.40-4.11) .001	3.54 (2.23-5.62) <.001	5.22 (2.08-13.10) <.001
3 apathy symptoms	4.28 (1.86-9.84) .001	3.74 (1.76-7.94) <.001	11.00 (2,88-41.97) <.001

*compared to no apathy symptoms

All models adjusted for depressive symptoms minus apathy (0-12) and cohort status

Results presented as: HR (95% CI) p