



Supplementary Material

Article Title: Antipsychotic Use and the Risk of Hip Fracture Among Community-Dwelling Persons With Alzheimer's Disease

Authors: Marjaana Koponen, MSc; Heidi Taipale, PhD; Piia Lavikainen, MSc; Antti Tanskanen, Phil Lic; Jari Tiihonen, MD, PhD; Anna-Maija Tolppanen, PhD; Riitta Ahonen, PhD; and Sirpa Hartikainen, MD, PhD

DOI Number: 10.4088/JCP.15m10458

List of Supplementary Material for the article

1. [eAppendix 1](#) Number of Hip Fractures Among Users of Individual Antipsychotic Drugs

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

eAppendix 1. Number of Hip Fractures among Users of Individual Antipsychotic Drugs

Antipsychotic drug	Number of users	Person years of use	Number of hip fractures
Risperidone	10,630	8933.4	272
Quetiapine	4,990	5343.9	143
Haloperidol	597	305.3	8
Olanzapine	249	232.2	9
Melperone	198	155.0	3
Perphenazine	127	107.9	4
Levomepromazine	57	43.4	0
Periciazine	39	29.2	1
Flupentixol	28	24.3	3
Sulpiride	23	15.5	0
Aripiprazole	10	6.9	0
Zuclopenthixol	7	2.4	1
Chlorprotixene	6	6.7	0
Chlorpromazine	5	1.7	0
Clozapine	3	2.0	0
Dixyrazine	2	0.3	0
Ziprasidone	1	0.5	0