

Supplementary Material

Article Title: Seasonal Variations of Australian Medicare-Reimbursed Psychiatric Consultations

Between 2016 and 2023: A Time Series Analysis

Author(s): Luke S.-C. Woon, DrPsych; David Smith, PhD; David F. Della, MD; Jeffrey C.L.

Looi, DMedSc; and Tarun Bastiampillai, MBBS

DOI

Number: https://doi.org/10.4088/PCC.24m03898

LIST OF SUPPLEMENTARY MATERIAL FOR THE ARTICLE

- Supplementary Table 1. MBS Items for Face-to-Face, Video, and Telephone Psychiatric Consultations
- 2. Supplementary Table 2. Sensitivity Analysis of Linear Regression Models for Quarterly Overall, Men's, and Women's Total Psychiatric Consultations With 3 Different Weighted Post-COVID Consultation Counts (85%, 80%, and 90%)
- 3. Supplementary Figure 1. Seasonal Plot of Total Quarterly MBS Psychiatric Consultations (2016-2023)
- 4. Supplementary Figure 2. Seasonal Plot of Quarterly MBS Psychiatric Telehealth Consultations (2016-2023)

DISCLAIMER

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary Table 1. MBS Items for Face-to-Face, Video, and Telephone Psychiatric Consultations

Service	Face-to-face	Video	Telephone
Consultation, not more than 15 minutes	300, 364*‡	91827, 353*†	91837
Consultation, 15 to 30 minutes	302, 366*‡	91828, 355*†	91838
Consultation, 30 to 45 minutes	304, 367*‡	91829, 356*†	91839
Consultation, 45 to 75 minutes	306, 369*‡	91830, 357*†	91840*
Consultation, more than 75 minutes	308, 370*‡	91831, 358*†	91841*
Prepare a treatment and management plan, patient under			
13 years with autism or another pervasive developmental	289	92434	92474*
disorder, at least 45 minutes			
Prepare a management plan, more than 45 minutes	291	92435	92475*
Review management plan, 30 to 45 minutes	293	92436, 359*†	92476*
Attendance, new patient (or has not received attendance i preceding 24 months), more than 45 minutes	n 296	92437, 361*†	92477*
Interview of a person other than patient, during initial diagnostic evaluation of patient, 20 to 45 minutes	348	92458	92498*
Interview of a person other than patient, during initial diagnostic evaluation of patient, 45 minutes or more	350	92459	92499*

Interview of a person other than patient, while continuing			
management of patient, not less than 20 minutes, not	352	92460	92500*
exceeding 4 attendances per calendar year			
Group psychotherapy, at least 1 hour, involving group of 2			
to 9 unrelated patients or a family group of more than 3	342	92455	92495*
patients, each referred to consultant psychiatrist			
Group psychotherapy, at least 1 hour, involving family			
group of 3 patients, each referred to consultant	344	92456	92496*
psychiatrist			
Group psychotherapy, at least 1 hour, involving family			
group of 2 patients, each referred to consultant	346	92457	92497*
group of 2 patients, each referred to consultant psychiatrist	346	92457	92497*
psychiatrist	346 90260	92457 92162, 90262*	92497* 92166*
psychiatrist Prepare an eating disorder treatment and management			
psychiatrist Prepare an eating disorder treatment and management plan, more than 45 minutes	90260	92162, 90262*	92166*

Notes: *Abolished items; †Telepsychiatry for patients in regional, remote, or rural areas; ‡ Face-to-face attendance following a telepsychiatry consultation.

Supplementary Table 2. Sensitivity Analysis of Linear Regression Models for Quarterly Overall, Men's, and Women's Total Psychiatric Consultations With 3 Different Weighted Post-COVID Consultation Counts (85%, 80%, and 90%). Constants are shown for Jan-Mar, while regression coefficients with their confidence intervals and p-values are shown for Apr-Jun, Jul-Sep, and Oct-Dec.


Quarter	Jan-Mar (referent)	Apr-Jun	Jul-Sep	Oct-Dec
85% post-COVID co	ounts			
Total	424578.5	441712.5	457408.7	437407.6
		[432555.9-	[448231.7-	[428196.7-
		450869.1]	466585.7]	446618.5]
		p=0.001*	p<0.001*	p=0.008*
Men	175394.2	181732.1	189238.2	180816.6
		[176356.7-	[183850.8-	[175409.3-
		187107.5]	194625.6]	186223.9]
		p=0.023*	p<0.001*	p=0.049*
Women	249184.3	259980.5	268170.5	256591.0
		[254755.3-	[262933.8-	[251334.9-
		265205.6]	273407.3]	261847.1]
		p<0.001*	p<0.001*	p=0.007*

80% post-COVID counts


Total	413589.9	428427.47	444667.58	426469.65
		[414147.1-	[430355.4-	[412104.6-
		442707.8]	458979.8]	440834.7]
		p=0.042*	p<0.001*	p=0.077
Men	170998.4	176399.0	184139.6	176459.7
		[168566.6-	[176289.7-	[168580.9-
		184231.4]	191989.4]	184338.6]
		p=0.169*	p=0.002*	p=0.166
Women	242591.5	252028.5	260528.0	250009.9
		[244924.6-	[253408.3-	[242863.9-
		259132.4]	267647.8]	257155.9]
		p=0.011*	p<0.001*	p=0.042*
90% post-CO	VID counts			
Total	435567.0	454997.5	470149.7	448345.5
		[447133.0-	[462267.7-	[440434.4-
		462862.0]	478031.7]	456256.6]
		p<0.001*	p<0.001*	p=0.003*
Men	179790.0	187,065	194,337	185,173
		[183502.5-	[190766.2-	[181589.7-
		190627.8]	197907.4]	188757.2]

		p<0.001*	p<0.001*	p=0.005*	
Women	255777.0	267932.3	275812.9	263172.0	
		[261713.4-	[269580.1-	[256916.2-	
		274151.3]	282045.7]	269427.8]	
		p<0.001*	p<0.001*	p=0.022*	

^{*}Statistically significant



Supplementary Figure 1. Seasonal Plot of Total Quarterly MBS Psychiatric Consultations (2016-2023)



Supplementary Figure 2. Seasonal Plot of Quarterly MBS Psychiatric Telehealth Consultations (2016-2023)